



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/29, 15/82, 5/10, A01H 5/00	A2	(11) International Publication Number: WO 99/53067 (43) International Publication Date: 21 October 1999 (21.10.99)
(21) International Application Number: PCT/CA99/00293 (22) International Filing Date: 13 April 1999 (13.04.99) (30) Priority Data: 09/059,090 13 April 1998 (13.04.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/059,090 (CIP) Filed on 13 April 1998 (13.04.98) (71) Applicant (for all designated States except US): HER MAJESTY IN RIGHT OF CANADA as represented by THE MINISTER OF AGRICULTURE AND AGRI-FOOD CANADA [CA/CA]; Eastern Cereal & Oilseed Research Centre, K.W. Neathy Building, Ottawa, Ontario K1A 0C6 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): MIKI, Brian [CA/CA]; 1876 Dorset Drive, Ottawa, Ontario K1H 5V1 (CA). GJZEN, Mark [CA/CA]; 848 Princess Avenue, London, Ontario N5W 3M4 (CA). MILLER, Shea [CA/CA]; 69 Kenilworth Street, Ottawa, Ontario K1Y 3Y5 (CA). BOW-	MAN, Lu-Ann [CA/CA]; R.R. #3, Listowel, Ontario N4W 3G8 (CA). BATCHELOR, Anthea [GB/CA]; 45 Alexander Street, Ottawa, Ontario K1M 1N1 (CA). HU, Ming [CA/CA]; 279 Kundson Drive, Kanata, Ontario K2K 2N8 (CA). BOUTILIER, Kim [CA/NL]; Groen van Prinsterstraat 87, NL-6702 CP Wageningen (NL). (74) Agents: SECHLEY, Konrad, A. et al.; Gowling, Strathy & Henderson, Suite 2600, 160 Elgin Street, Ottawa, Ontario K1P 1C3 (CA). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: SEED-COAT PROMOTERS, GENES AND GENE PRODUCTS (57) Abstract <p>The present invention is directed to isolated genomic sequences that are differentially expressed within seed-coat tissues. These DNAs are expressed within the inner integument, thick, or thin, walled parenchyma, endothelium, palisade, hourglass, or stellate parenchyma cells of the seed-coat. Furthermore, this invention relates to promoter regions obtained from genomic sequences that are differentially expressed in seed-coat tissues, and their use for directing seed-coat specific expression of genes of interest within transformed plant cells or plants. Exemplified promoters include those obtained from the differential screening of a seed-coat library and cryptic promoters obtained using T-DNA tagging using a promoterless marker gene.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

SEED-COAT PROMOTERS, GENES AND GENE PRODUCTS

Field of Invention

5 This invention relates to seed-coat promoters, genes and proteins encoded by these genes. More specifically, it relates to genes and promoters that are developmentally regulated and expressed, or activated, within tissues comprising the seed-coat, and tissues directly associated with the seed-coat, of plants. Furthermore, this invention also relates to proteins encoded by genes
10 expressed within these tissues are their localization within, or onto, the seed-coat.

Background and Prior Art

15 Bacteria from the genus *Agrobacterium* have the ability to transfer specific segments of DNA (T-DNA) to plant cells, where they stably integrate into the nuclear chromosomes. Analyses of plants harbouring the T-DNA have revealed that this genetic element may be integrated at numerous locations, and can occasionally be found within genes. One strategy which may be exploited
20 to identify integration events within genes is to transform plant cells with specially designed T-DNA vectors which contain a reporter gene, devoid of *cis*-acting transcriptional and translational expression signals (i.e. promoterless), located at the end of the T-DNA. Upon integration, the initiation codon of the promoterless gene (reporter gene) will be juxtaposed to plant sequences. The
25 consequence of T-DNA insertion adjacent to, and downstream of, gene promoter elements may be the activation of reporter gene expression. The resulting hybrid genes, referred to as T-DNA-mediated gene fusions, consist of unknown and thus uncharacterized plant promoters residing at their natural location within the chromosome, and the coding sequence of a marker gene
30 located on the inserted T-DNA (Fobert *et al.*, 1991, Plant Mol. Biol. 17, 837-851).

-2-

It has generally been assumed that activation of promoterless or enhancerless marker genes result from T-DNA insertions within or immediately adjacent to genes. The recent isolation of several T-DNA insertional mutants (Koncz *et al.*, 1992, *Plant Mol. Biol.* **20**, 963-976; reviewed in Feldmann, 1991, *Plant J.* **1**, 71-82; Van Lijsebettens *et al.*, 1991, *Plant Sci.* **80**, 27-37; Walden *et al.*, 1991, *Plant J.* **1**: 281-288; Yanofsky *et al.*, 1990, *Nature* **346**, 35-39), shows that this is the case for at least some insertions. However, other possibilities exist. One of these is that integration of the T-DNA activates silent regulatory sequences that are not associated with genes. Lindsey *et al.* (1993, *Transgenic Res.* **2**, 33-47) referred to such sequences as "pseudo-promoters" and suggested that they may be responsible for activating marker genes in some transgenic lines.

Inactive regulatory sequences that are buried in the genome but with the capability of being functional when positioned adjacent to genes have been described in a variety of organisms, where they have been called "cryptic promoters" (Al-Shawi *et al.*, 1991, *Mol. Cell. Biol.* **11**, 4207-4216; Fourel *et al.*, 1992, *Mol. Cell. Biol.* **12**, 5336-5344; Irniger *et al.*, 1992, *Nucleic Acids Res.* **20**, 4733-4739; Takahashi *et al.*, 1991, *Jpn J. Cancer Res.* **82**, 1239-1244). Cryptic promoters can be found in the introns of genes, such as those encoding for yeast actin (Irniger *et al.*, 1992, *Nucleic Acids Res.* **20**, 4733-4739), and a mammalian melanoma-associated antigen (Takahashi *et al.*, 1991, *Jpn J. Cancer Res.* **82**, 1239-1244). It has been suggested that the cryptic promoter of the yeast actin gene may be a relict of a promoter that was at one time active but lost function once the coding region was assimilated into the exon-intron structure of the present-day gene (Irniger *et al.*, 1992, *Nucleic Acids Res.* **20**, 4733-4739). A cryptic promoter has also been found in an untranslated region of the second exon of the woodchuck N-myc proto-oncogene (Fourel *et al.*, 1992, *Mol. Cell. Biol.* **12**, 5336-5344). This cryptic promoter is responsible for activation of a N-myc2, a functional processed gene which arose from retroposition of N-myc transcript (Fourel *et al.*, 1992, *Mol.*

-3-

Cell. Biol. 12, 5336-5344). These types of regulatory sequences have not yet been isolated from plants.

5 Weber et al. (1995, *Plant Cell* 7:1835-1846) disclose a cDNA sequence of a seed-coat associated invertase. However, all of the cDNA's characterized were found to be expressed in tissues other than the seed-coat, including anthers, cotyledon, stem and root. Furthermore, no promoter was isolated, characterized, or disclosed.

10 Described herein is the occurrence of seed-coat genes and promoters that have been obtained as a result of differential screening of seed-coat genomic libraries, or generated by tagging with a promoterless GUS (β -glucuronidase) T-DNA vector, or by identification of genes that are highly
15 expressed in the seed-coat or associated tissues. Expression analysis of these DNA's reveal that they are spatially and developmentally regulated in seed coats. Prior to this work, promoters, as well as genes specifically expressed in seed coat tissues had not been isolated or reported. Furthermore, proteins encoded by genes that are expressed within seed-coat, or associated with seed-
20 coat tissues, are also disclosed.

Summary of Invention

25 This invention relates to seed-coat promoters and genes. More specifically, it relates to genes and promoters that are developmentally regulated and expressed, or activated, within tissues comprising the seed-coat of plants, and tissues directly associated with the seed-coat, of plants. Furthermore, this invention also relates to proteins encoded by genes expressed within these tissues and their localization within, or onto, the seed-coat..

30

-4-

A transgenic tobacco plant, T218, contained a 4.7 kb *EcoRI* fragment containing the 2.2 kb promoterless GUS-*nos* gene and 2.5 kb of 5' flanking tobacco DNA. Deletion of the region approximately between 2.5 and 1.0 kb of the 5' flanking region did not alter GUS expression, as compared to the entire 4.7 kb GUS fusion. A further deletion to 0.5 kb of the 5' flanking site resulted in complete loss of GUS activity. Thus the region between 1.0 and 0.5 of the 5' flanking region of the tobacco DNA contains the elements essential to gene activation. This region is contained within a *XbaI* - *SnaBI* restriction site fragment of the flanking tobacco DNA. Furthermore, other promoters have been identified that are differentially expressed within the seed-coats of plants, and that are capable of driving expression of heterologous genes that are operatively linked thereto.

Thus according to the present invention there is provided an isolated genomic DNA molecule, differentially expressed in seed coat tissues. Furthermore, this genomic DNA molecule is differentially expressed within the outer integument of the seed coat, the inner integument of the seed coat, the thick walled parenchyma of the seed coat, the thin walled parenchyma of the seed coat, the endothelium of the seed coat, the hourglass cells of the seed coat, the palisade of the seed coat, the stellate parenchyma of the seed coat, or the membranous endocarp, or a combination thereof.

This invention is also directed to a seed-coat promoter obtained from the genomic DNA molecule as described above. Also considered within the scope of the present invention is a cryptic seed coat promoter. Furthermore, this invention is directed to a seed coat promoter, as described above, that controls the differential expression of a gene associated therewith, within the outer integument of the seed coat, the inner integument of the seed coat, the thick walled parenchyma of the seed coat, the thin walled parenchyma of the seed coat, the endothelium of the seed coat, the hourglass cells of the seed coat, the

-5-

palisade of the seed coat, or the stellate parenchyma the seed coat, membranous endocarp, or a combination thereof

5 This invention also relates to an isolated genomic DNA characterized by the restriction map selected from the group consisting of Figure 12 (a), Figure 12 (b), Figure 12 (c) and Figure 12 (d).

10 According to the present invention, there is also provided an isolated seed-coat promoter. Furthermore, this seed coat promoter may be obtained from angiosperms. More specifically, this seed-coat promoter is obtained from the group consisting of tobacco or soybean.

15 This invention is also directed to a cloning vector comprising a gene encoding a protein and an isolated seed-coat promoter, wherein the gene is under the control of the seed-coat promoter. Furthermore, this invention includes a plant cell which has been transformed with such a vector.

20 This invention also provides for a transgenic plant containing a seed-coat promoter, operatively linked to a gene encoding a protein.

The present invention is also directed to a seed-coat promoter comprising at least 10 contiguous nucleotides of nucleotides 1-2526 of SEQ ID NO:7, or an analogue of the sequence defined by nucleotides 1-2526 of SEQ ID NO:7, wherein the analogue hybridizes to a nucleic acid defined by nucleotides 1-2526 of SEQ ID NO:7 under stringent hybridization conditions and maintains seed-coat, or seed-coat associated promoter activity.

30 This invention also includes a seed-coat promoter comprising at least 10 contiguous nucleotides of nucleotides 1-2450 of SEQ ID NO:8, or an analogue of the nucleic acid sequence defined by nucleotides 1-2450 of SEQ ID NO:8, wherein the analogue hybridizes to a nucleic acid defined by nucleotides

-6-

1-2450 of SEQ ID NO:8 under stringent hybridization conditions and maintains seed-coat, or seed-coat associated promoter activity.

The present invention also is directed to a seed-coat promoter comprising at least 10 contiguous nucleotides of nucleotides 1-5514 of SEQ ID NO:9, or an analogue of the nucleotides sequence defined by nucleotides 1-5514 of SEQ ID NO:9, wherein the analogue hybridizes to a nucleic acid defined by nucleotides 1-5514 of SEQ ID NO:9 under stringent hybridization conditions and maintains seed-coat, or seed-coat associated promoter activity.

10

Brief Description of the Drawings

Figure 1 depicts the fluorogenic analyses of GUS expression in the plant T218. Each bar represents the average \pm one standard deviation of three samples. Nine different tissues were analyzed: leaf (L), stem (S), root (R), anther (A), petal (P), ovary (O), sepal (Se), seeds 10 days post anthesis (S1) and seeds 20 days post-anthesis (S2). For all measurements of GUS activity, the fraction attributed to intrinsic fluorescence, as determined by analysis of untransformed tissues, is shaded black on the graph. Absence of a black area at the bottom of a histogram indicates that the relative contribution of the background fluorescence is too small to be apparent.

Figure 2 shows the cloning of the GUS fusion in plant T218 (pT218) and construction of transformation vectors. Plant DNA is indicated by the solid line and the promoterless GUS-*nos* gene is indicated by the open box. The transcriptional start site and presumptive TATA box are located by the closed and open arrow heads respectively. DNA probes #1, 2, 3 and RNA probe #4 are shown. The *EcoRI* fragment in pT218 was subcloned in the pBIN19 polylinker to create pT218-1. Fragments truncated at the *XbaI* *SnaBI* and *XbaI* sites were also subcloned to create pT218-2, pT218-3 and pT218-4.

-7-

Abbreviations for the endonuclease restriction sites are as follows: *Eco*RI (E), *Hind*III (H), *Xba*I (X), *Sna*BI (N), *Sma*I (M), *Sst*I (S).

Figure 3 shows the expression pattern of promoter fusions during seed development. GUS activity in developing seeds (4-20 days postanthesis (dpa)) of (Fig. 3a) plant T218 (●-●) and (Fig. 3b) plants transformed with vectors pT218-1 (○-○), pT218-2 (□-□), pT218-3 (▽-▽) and pT218-4 (Δ-Δ) which are illustrated in Figure 2. The 2 day delay in the peak of GUS activity during seed development, seen with the pT218-2 transformant, likely reflects greenhouse variation conditions.

Figure 4 shows GUS activity in 12 dpa seeds of independent transformants produced with vectors pT218-1 (○), pT218-2 (□), pT218-3 (▽) and pT218-4 (Δ). The solid markers indicate the plants shown in Figure 3 (b) and the arrows indicate the average values for plants transformed with pT218-1 or pT218-2.

Figure 5 shows the mapping of the T218 GUS fusion termini and expression of the region surrounding the insertion site in untransformed plants. Figure 5(a) shows the mapping of the GUS mRNA termini in plant T218. The antisense RNA probe from subclone #4 (Figure 2) was used for hybridization with total RNA of tissues from untransformed plants (10 μg) and from plant T218 (30 μg). Arrowheads indicate the anticipated position of protected fragments if transcripts were initiated at the same sites as the T218 GUS fusion. Figure 5 (b) shows the RNase protection assay using the antisense (relative to the orientation of the GUS coding region) RNA probe from subclone e (Figure 7) against 30 μg total RNA of tissues from untransformed plants. P, untreated RNA probe; -, control assay using the probe and tRNA only; L, leaves from untransformed plants; 8, 10, 12, seeds from untransformed plants at 8, 10, and 12 dpa, respectively; T10, seeds of plant T218 at 10 dpa; +, control hybridization against unlabelled *in vitro*-synthesized sense RNA from subclone

-8-

c (panel a) or subclone e (panel b). The two hybridizing bands near the top of the gel are end-labelled DNA fragment of 3313 and 1049 bp, included in all assays to monitor losses during processing. Molecular weight markers are in number of bases.

5

Figure 6 provides the nucleotide sequence of pT218 (top line) (SEQ ID NO: 1) and pIS-1 (bottom line). Sequence identity is indicated by dashed lines. The T-DNA insertion site is indicated by a vertical line after bp 993. This site on pT218 is immediately followed by a 12 bp filler DNA, which is followed by the T-DNA. The first nine amino acids of the GUS gene and the GUS initiation codon (*) are shown. The major and minor transcriptional start site is indicated by a large and small arrow, respectively. The presumptive TATA box is identified and is in boldface. Additional putative TATA and CAAT boxes are marked with boxes. The location of direct (1-5) and indirect (6-8) repeats are indicated by arrows.

10

15

Figure 7 shows the base composition of region surrounding the T218 insertion site cloned from untransformed plants. The site of T-DNA insertion in plant T218 is indicated by the vertical arrow. The position of the 2 genomic clones pIS-1 and pIS-2, and of the various RNA probes (a-e) used in RNase protection assays are indicated beneath the graph.

20

Figure 8 shows the Southern blot analyses of the insertion site in *Nicotiana* species. DNA from *N. tomentosiformis* (N tom), *N. sylvestris* (N syl), and *N. tabacum* (N tab) were digested with *Hind*III (H), *Xba*I (X) and *Eco*RI (E) and hybridized using probe #2 (Figure 2). Lambda *Hind*III markers (kb) are indicated.

25

Figure 9 shows the AT content of 5' non-coding regions of plant genes. A program was written in PASCAL to scan GenBank release 75.0 and to calculate the AT contents of the 5' non-coding (solid bars) and the coding

30

regions (hatched bars) of all plant genes identified as "Magnoliophyta" (flowering plants). The region -200 to -1 and +1 to +200 were compared. Shorter sequences were also accepted if they were at least 190 bp long. The horizontal axis shows the ratio of the AT content (%). The vertical axis shows the number of the sequences having the specified AT content ratios

Figure 10 shows a Northern analysis of the expression of several of the genes of the present invention within developing seed coats, embryo, pod, flower, root, stem and leaf tissues. Figure 10 (a) shows the expression of SC4; Figure 10 (b) shows the expression of SC20; Figure 10 (c) shows the expression of SC21, Figure 10 (d) shows the expression of *Ep* locus peroxidase within these tissues. Figure 10 (e) shows the expression of HP (hydrophobic protein) in leaf, flower, pod, seed coat, embryo, stem or root tissues. Figures 10 (f) and (g), total RNA was isolated from leaf, flower, pod shells, seed coat, embryo, stem or root tissue. Equal amounts of RNA (10 μ g) were vacuum blotted to nylon and probed with *HPS* cDNA. Ribosomal RNA (rRNA), visualized by staining with ethidium bromide, is shown as control. Figure 10 (f), RNA from tissues at early (E) mid (M) or late (L) stages of development were compared for *HP* gene expression. All samples shown are from dull seeded phenotype (cv Harosoy 63). Figure 10 (g), RNA from pod tissues of dull (cv Harosoy 63) and shiny (cv. Williams 82) seeded soybeans were compared for *HP* gene expression.

Figure 11 shows the restriction maps obtained from Figure 11 (a) SC20; Figure 11 (b) SC21; Figure 11 (c) HP (hydrophobic protein) genomic region, and Figure 11 (d) SC4. Included in Figure 11 (c) are restriction enzyme sites for *Bam*HI, *Bgl*II, *Hind*III, and *Xba*I; the HP ORF; TATA box consensus signals; and the position of direct repeats of 12 bp or longer.

-10-

Figure 12 shows the morphology of the seed coat of *Glycine max*.

Figure 12 (a) shows the structures present at six days after anthesis (DAF);

Figure 12 (b), at 12 DAF; and Figure 12 (c) at 18 DAF.

5 Figure 13 shows *in situ* hybridization results obtained with seed coats of
Glycine max at different developmental stages, and probed as follows: Figure
 13 (a) seed coat at 3 days after anthesis (DAF), probed with SC4; Figure 13
 (b) seed coat at 9 DAF, probed with SC20; Figure 13 (c) seed coat at 15 DAF,
 probed with SC21; Figure 13 (d) seed coat at 18 days after anthesis, probed
 10 with a soybean peroxidase, corresponding to the *Ep* locus. Figures 13 (e), (f)
 and (g) were obtained from cross sections of developing soybean seeds (cultivar
 Maple Presto, *EpEp*). Hybridization of ^{35}S -probe to complementary mRNA
 appears as bright white signal in these dark field microscopy images. Figure
 13 (e) 6 DAF (DPA, days post anthesis), Figure 13 (f) 9 DAF, and Figure 13
 15 (g) 12 DAF. Scale bars are 100 μm . Emb, embryo; F, funiculus; HG,
 hourglass cells; PC, pericarp; SC, seed coat.

Figure 14 shows light micrographs of a seed-coat obtained from
 soybean. Figure 14 (a) shows a plastic embedded section of the seed-coat near
 20 the hilum at 21 daf and stained with Toluidine Blue O. Note the association of
 the membranous endocarp with the seed-coat pallisade. Figure 14 (b) shows a
 wax-embedded section of a soybean seed-coat as 12 daf probed with ^{35}S -
 labelled Hydrophobic Protein (HP) antisense RNA, and counter stained with
 Toluidine Blue O. Note strong specific localization of the probe within the
 25 membranous endocarp. Pallisade (p), hourglass cells (h), counterpallisade (c),
 arial cells (a), stellate parenchyma (s), thin walled parenchyma (n), thickwalled
 parenchyma (k), pod parenchyma (d), and membranous endocarp ►. Figures
 14 (c) and (d). show localization of *HP* mRNA transcript by *in situ*
 hybridization. Cross sections of soybean pods containing immature seeds (dull
 30 phenotype, HPS (+), cv Maple Presto). Hybridization of ^{35}S labelled *HP* probe
 to complementary mRNA appears as bright white signal in these dark field

microscopy images. E, embryo; Ep, inner epidermal layer of endocarp; Ex, exocarp; F, funiculus; M, mesocarp; Sc, seed coat; Sm, sclerenchyma layer of endocarp. Bar = 100 μ m. Figure 14 (c), Expression at 6 DPA (days post anthesis). Figure 14 (d), Expression 12 DPA.

5

Figure 15 shows the Soybean hydrophobic protein (HP) cDNA and deduced amino acid sequences. Figure 15 (a), the cDNA and amino acid sequence of HP. The pre-protein signal sequence is underlined. Figure 15 (b) shows the deduced amino acid sequence of HP pre-protein. Alternate N-terminal residues are boxed, as determined by peptide microsequence analysis. Figure 15 (c) shows a Kyle-Doolittle hydrophilicity plot of HP (Lasergene). In this plot, positive values indicate greater hydrophilic character. Also represented are the three domains of the HP pre-protein and the length of the mature peptide. Figure 15 (d) shows a schematic comparison of HP domain structure to three other plant proteins. Bold numbers indicate the length in amino acid residues for the domain segments. The pattern of spacing between the eight cysteine residues within the hydrophobic domains is also shown below each protein. Sequences for tobacco N16 polypeptide (D86629), maize proline rich hydrophobic protein (PRHP) (X60432), and *Arabidopsis* lipid transfer protein 1 (LTP1) (M80567) were retrieved from GenBank.

10
15
20

Figure 16 shows scanning electron micrographs of representative 'Dull' and 'Shiny' seeded soybean cultivars. Scale bars are included in the figures. The lowest magnification (x18), Figure 16 (a) is a view of the entire seed. The large oval shaped scar on the seed surface is the hilum, corresponding to the point of detachment of the mature seed from the funiculus. Figure 16 (b), x100, and Figure 16 (c) x500, are focused outside of hilum region.

25

Figure 17 shows a silver stained SDS-PAGE analysis of protein extracts from seed tissues and surface. Lanes marked 'M' indicate protein standards, and their corresponding mass in kilodaltons is also provided. Figure 17 (a),

30

-12-

Soluble protein extracts from the embryo, seed coat, and seed surface of a dull phenotype (cv Harosoy 63). Each sample at approximately 1 μ g of total protein. Figure 17 (b), Seed surface protein extracts of a dull phenotype (cv Harosoy 63) with different concentrations of dithiothreitol (DTT) present in the sample loading buffer, as indicated at the top of each lane. Figure 17 (c), Seed surface protein extracts of dull (D), shiny (S), and bloom (B).

Figure 18 shows restriction fragment length polymorphisms between dull and shiny phenotypes. Genomic DNA from dull (cv Harosoy 63) and shiny (cv Williams 82) soybeans with abundant (+) or trace (-) amounts of HPS on the seed surface, was digested with restriction enzymes, electrophoretically separated, blotted, and hybridized to HP cDNA probe. The size of hybridizing fragments was estimated by comparison with standards and is shown on the left.

15

Figure 19 shows the nucleotide sequence and deduced amino acid sequence of SC4 cDNA, and the sequence comparisons between SC4 protein and BURP proteins. Figure 19 (a), 5' and 3' untranslated sequences are in lowercase lettering. The stop codon is shown with an asterisk and two polyadenylation signals are underlined. Two copies of a ten amino acid repeat is also underlined. Consensus sequences for N-glycosylation (NNT; NSSN; and NGTV) are also underlined. Figure 19 (b), amino acid alignment of the carboxyl terminus of the SC4 protein with the BURP domain (A) and the amino terminus of the SC4 protein with the conserved segments of the second domain (B) of several BURP domain proteins. Pg1 β is not included in panel B as the second domain of this protein does not contain a conserved segment. Gaps were introduced to optimize the alignment. Conserved amino acids are shown in bold face. Amino acids of each protease are numbered from the precursor sequence. Figure 19 (c) shows the structural similarity between SC4 protein and the BURP domain proteins.

30

-13-

Figure 20 shows Northern blot analysis of SC4 and SC20 mRNA accumulation in seed coat embryo and pod organs of soybean. 10 μ g total RNA from seed coat, embryo and pod organs between 6-24 days past anthesis were hybridized with radiolabelled probes. For day 6, total RNA was prepared from whole seeds. Each blot was hybridized with a SC4 cDNA probe, Figure 20 (a), a SC20 cDNA probe Figure 20 (b), and an 18S rRNA probe Figure (c).

Figure 21, shows the localization of SC4 mRNA in Seed coat organs of soybean by *in situ* hybridization. Transections of seed coats at 3 days past anthesis (dpa) and 6 dpa. Hybridization to Antisense, Figure 21 (a), and sense, Figure 21 (b) SC4 labelled RNA probes. Abbreviations, II - inner integument, OI outer integument, P pod. Bar equals 100 μ m.

Figure 22, shows Southern blot analysis of SC4. Figure 22 (a) shows Southern analysis of the gene family composition of *sc4* in soybean. Figure 22 (b) shows Southern analysis of *sc4* in diverse plant species. Hybridized filter was washed under conditions of low stringency, twice at 52°C for 15 min in 2x SSC, 0.1% SSC, 0.1% SDS and once at 52°C for 30 min in 0.1x SSC, 0.1% SDS.

Figure 23 reveals the characterization of *sc20* and the SC20 protein. Figure 23 (a) is a restriction map of *sc20*. Figure 23 (b) shows the nucleotide sequence and deduced amino acid sequence of *sc20* cDNA. The stop codon is shown with an asterisk and the polyadenylation signal is underlined. The consensus sequences for N-glycosylation are also underlined. Figure 23 (c) shows the hydrophobic plot of SC20 protein, where hydrophobic regions possess a positive sign, and hydrophilic regions possess a negative sign. In Figure 23 (d), alignment of SC20 protein with other subtilases is shown. D, H and S regions represent amino acid sequences around the catalytic aspartate, histidine and serine residues of the subtilases. The catalytic residues are labelled with an asterisk. N region represents amino acid sequence around the

-14-

conserved asparagine residue, of subtilases. # indicates the conserved asparagine. AF70, cucumisin, P69B, Ag12, subtilisin BPN', kex2, furin are from *Picea abies*, *Cucumis melo* L., *Lycopersicon esculentum*, *Alnus glutinosa*, *Bacillus subtilis*, *Saccharomyces cerevisiae*, and *Homo sapiens* respectively.

5 Conserved amino acids are shown in boldface. Amino acids of each protease are numbered from the precursor sequence.

Figure 24 shows localization of SC20 mRNA in seed coats of soybean by in situ hybridization. Transection of seed coats at 12 days past anthesis hybridized to Antisense, Figure 24 (a), and Sense, figure 24 (b), SC20 radiolabelled RNA probes. Abbreviations: H Hilumi, II inner integument, OI outer integument, * thick walled parenchyma, ** thin walled parenchyma. Bar equals 100µm.

15 Figure 25 shows Southern blot analysis of *sc20*. Figure 25 (a) and (b), Southern analysis of the gene family composition of *sc20* in soybean under conditions of medium stringency (twice at 52°C for 15 min in 2x SSC, 0.1% SDS, and once at 52°C for 30 min in 0.1x SSC, 0.1% SDS), Figure 25 (a), and high stringency (once at 62°C for 30 min in 0.1x SSC, 0.1% SDS) Figure 25 (b). Figure 25 (c) shows Southern analysis of *sc4* in diverse plant species. genomic DNA was digested with EcoRI. Hybridization used a radiolabelled SC20 cDNA probe. The filter was washed under conditions of medium stringency, twice at 52°C for 15 min in 2x SSC, 0.1% SDS and once at 52°C for 30 min in 0.1x SSC, 0.1% SDS.

25

Detailed Description of the Preferred Embodiments

T-DNA tagging with a promoterless β -glucuronidase (GUS) gene generated a transgenic *Nicotiana tabacum* plant that expressed GUS activity only in developing seed coats. Cloning and deletion analysis of the GUS fusion

30

-15-

revealed that the promoter responsible for seed coat specificity was located in the plant DNA proximal to the GUS gene. Deletion analyses localized the cryptic promoter to an approximately 0.5 kb region between a *Xba*I and a *Sna*BI restriction endonuclease site of the 5' flanking tobacco DNA. This region spans from nucleotide 1 to nucleotide 467 of SEQ ID NO: 1.

Other work, based on the differential screening of seed coat libraries has led to the identification of several other genes that are differentially expressed within, or tissues that are directly associated with, the seed coat of plants. These genes include SC4 (SEQ ID NO's: 3 and 9, cDNA and genomic sequences, respectively), SC20 (SEQ ID NO's: 4 and 8, cDNA and genomic sequences respectively), SC21 (SEQ ID NO: 5, cDNA sequence), and their associated promoters (see SEQ ID NO 9 and 8 for promoters of SC 4 and SC20, respectively; also Figure 12). Furthermore, the isolation of genes encoding highly expressed seed coat proteins led to the identification of a seed-coat specific peroxidase from the *Ep* locus and associated promoter (*Ep* genomic sequence, SEQ ID NO:2), as well as a gene encoding a seed-coat localized hydrophobic protein (HP, cDNA sequence SEQ ID NO:6) and associated promoter (within genomic sequence, SEQ ID NO:7, also see Figure 11 (c)). Thus, the present invention includes promoters, genes and proteins isolated from several plant species, that are preferentially expressed, or specific to seed-coat tissues, as well as promoters, genes and associated proteins obtained from tissues associated with the seed-coat.

The term cryptic promoter means a promoter that is not associated with a gene and thus does not control expression in its native location. These inactive regulatory sequences are buried in the genome but are capable of being functional when positioned adjacent to a gene.

The DNA sequence of an aspect of the present invention includes the DNA sequence of SEQ ID NO: 1, the promoter region within SEQ ID NO: 1

-16-

(for example from nucleotide 1 to 476), and analogues thereof. Similarly, another aspect of this invention includes a DNA sequence of SEQ ID NO:2, the promoter region of this sequence (nucleotides 1-1532), and analogues thereof. Another aspect of this invention includes a DNA sequence of SEQ ID NO:7, the promoter region (nucleotides 1-2526), and analogues thereof, a DNA sequence of SEQ ID NO 8, the promoter region (nucleotides 1-2450) and analogues thereof, and a DNA sequence of SEQ ID NO:9, the promoter region (nucleotides 1-5514) and analogues thereof.

10 Analogues include those DNA sequences which hybridize under stringent hybridization conditions (see Maniatis *et al.*, in Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory, 1982, p. 387-389) to the DNA sequence of SEQ ID NO: 1, 2, 7, 8 or 9 provided that said sequences maintain the seed coat, or seed-coat associated promoter activity. An example of one such stringent hybridization conditions may be hybridization at 4XSSC at 65°C, followed by washing in 0.1XSSC at 65°C for an hour, or at 62°C for 30 min in 0.1x SSC, 0.1% SDS. Alternatively an exemplary stringent hybridization condition could be in 50% formamide, 4XSSC at 42°C. With the use of Digoxigenin labelled probes, stringent hybridization may include 65 °C in 0.25 M Na₂HPO₄ (pH 7.2), 20% SDS, 1 mM EDTA and 0.5% blocking reagent (Boehringer Mannheim) followed by washing at 22 °C in 20 mM Na₂HPO₄ (pH 7.2), 1% SDS and 1 mM EDTA and washes in the same solution at 68 °C. Analogues also include those DNA sequences which hybridize to the sequence of SEQ ID NO: 1, 2, 7, 8 or 9 under relaxed hybridization conditions provided that said sequences maintain the seed-coat promoter activity. Examples of such non-hybridization conditions includes hybridization at 4XSSC at 50°C or with 30-40% formamide at 42°C. Alternate conditions of medium stringency include washing the filter twice at 52°C for 15 min in 2x SSC, 0.1% SDS and once at 52°C for 30 min in 0.1x SSC, 0.1% SDS.

30

-17-

Furthermore, another aspect of this invention is directed to the identification and characterization of seed-coat promoters (see Figure 11) and their corresponding genes of cDNA's (SEQ ID NO's: 3-6), as characterized by Southern or *in situ* hybridization analysis of the expression patterns of genes expressed under the control of seed-coat promoters within developing seed coats (Figures 13 and 14). Furthermore, restriction maps of the promoter and downstream regions of several seed-coat genomic clones is presented (Figure 11).

Proteins of interest may be expressed in seed coat tissues by placing a gene capable of expressing the protein of interest under the control of the DNA promoters of this invention. Genes of interest include but are not restricted to herbicide resistant genes, genes encoding viral coat proteins, or genes encoding proteins conferring biological control of pest or pathogens such as an insecticidal protein for example *B. thuringiensis* toxin. Other genes include those capable of modifying the production of proteins that alter the taste of the seed and/or that affect the nutritive value of the seed.

By "seed-coat" it is meant tissues typically found within, and associated with, the seed-coat of developing or mature angiosperm seeds. Without wishing to limit the types of tissues found within a seed-coat, this region of the seed typically comprises a range of cell types including, and bounded by, an inner endothelium, and an outer epidermis or palisade cell layer. Within these inner and out cell layers, there may be found parenchyma-like cells, for example thin, or thick walled parenchyma, or stellate parenchyma, vascular tissue, hypodermis, hour-glass cells (osteosclereids), and one or more integuments, including the inner and outer integuments. However, it is to be understood that other cell types found within the region between the inner endothelium and outer epidermis may also be considered to comprise the seed-coat, for example, but not limited to the hilum, and funiculus comprising arial cells. Furthermore, it is to be understood that "seed-coat" also refers to tissues

-18-

associated with, or adhering to the seed coat, for example the membranous endocarp (of the inner ovary wall), as this cell type adheres to the seed-coat and remains in association with the seed coat (see for example Figure 15 (a), (b), and Figure 16). Therefore, as used herein, tissues that are associated with, or that adhere to, the seed-coat are referred to as "seed-coat associated tissues", or "tissues associated with the seed-coat". It is contemplated that other cell types may also associate with seed-coat tissues in addition to those disclosed above and that the tissues identified above should not be considered limiting in any manner.

By "seed-coat gene" it is meant a gene that is differentially expressed within the seed-coat as detected under stringent conditions (as defined above). Examples of such a gene include, but are not limited to SC4, SC 20, SC 21, or *Ep* locus peroxidase. However, the product of the gene may be exported from the cell to an exterior location of a seed-coat cell, including the surface of the seed-coat itself. An example, which is not to be considered limiting in any manner, of a gene product that is synthesized in within a seed-coat associated cell, and that is localized onto the surface of the seed coat, is the hydrophobic protein (HP; see Figures 14 (a) (b) and Figure 16).

A "seed-coat promoter" is a promoter that is differentially active within cells of the seed-coat. When operably linked with a gene under its control, a seed-coat promoter confers expression to a gene within the seed-coat, which can be detected under stringent conditions (as defined above). Seed coat associated promoter refers to a promoter that is active in tissue associated with the seed coat as defined above.

By "differentially expressed" it is meant the expression of a gene under the control of a promoter, as detected by standard means, within a specified tissue or organ. Such standard means for detecting expression include, but are not limited to, Northern, or *in situ* hybridizations and the like performed under

-19-

stringent conditions, or reporter gene expression. For example, a gene that is differentially expressed in seed-coat tissues is detectable within seed-coat tissues, and displays little or no expression in other tissues such as root, stem.

5 By "preferentially expressed" it is meant the expression of a gene under the control of a promoter, as detected by standard means, wherein the majority of expression is detected within a specified tissue or organ. Such standard means for detecting expression include, but are not limited to, Northern, or *in situ* hybridizations and the like performed under stringent conditions, or the
10 expression of reporter genes. For example, a gene that is preferentially expressed in seed-coat tissues is detectable within seed-coat tissues, but may exhibit some expression within other tissues such as root, stem.

By "seed-coat localized" or "localized onto the seed-coat" it is meant a
15 gene product that, as a result of some property of the amino acid sequence of the gene product, is targeted within, or onto seed-coat tissues, respectively. Properties of an amino acid sequence that may direct the targeting of a protein within or onto seed-coat tissues include, but are not limited to, signal sequences that direct intracellular, and extracellular localization of a protein, and also
20 hydrophobic regions within a protein, that results in localization of the protein onto the seed coat. An example, which is not to be considered limiting, of a protein that is localized onto the seed-coat, is the hydrophobic protein (HP). HP is localized on the outside of the seed-coat following its synthesis within the membranous endocarp, and appears to be involved with the adherence of the
25 endocarp to the seed-coat (see Figure 16).

Development of the Soybean coat

30 The seed coat of *Glycine max* undergoes dramatic changes in the first two and a half weeks after anthesis (flowering). At six days after anthesis (DAF; see Figure 12 (a)), the seed coat has a distinct epidermis (10), consisting

-20-

of thin-walled cuboidal cells; an outer integument (20), consisting of up to a dozen layers of thin-walled parenchyma, and containing vascular tissue (recurrent vascular bundle) in the subhilar region; an inner integument (30), consisting of up to 6 layers of deeply-staining thick-walled parenchyma; and an
5 endothelium (40), consisting of thin-walled cuboidal cells.

At 12 days after anthesis (Figure 12 (b)), there is a distinct hypodermis (15) of thin-walled cuboidal cells directly beneath the epidermis (10); the outer integument (20) has differentiated into an upper layer of thin-walled
10 parenchyma (25), and a lower layer of thick-walled parenchyma (27); the inner integument (30), while still having very thick, deeply staining cell walls, has become stretched, and is compressed to about 3 cells thick; the endothelium (40) is also retained. Also evident in Figure 12 (b) is the endosperm (50), and the developing cotyledons (60).

15 By 18 days after anthesis (Figure 12 (c)), the epidermal cells have divided and elongated to form thick-walled macrosclereids, forming a palisade layer (13). The hypodermis has differentiated into osteosclereids: thick walled cells with a characteristic I-shape (hourglass cells; 17). A prominent vascular
20 region (70) has developed in the thin-walled parenchyma (25) of the outer integument which stops before reaching the region of the seed opposite the hilum; the thick-walled parenchyma (27) is retained. The inner integument (30) has become completely stretched and crushed, leaving a single, deeply staining wall layer directly above the endothelium (40). The hilum region
25 contains a well-developed counter-palisade (80), and a tracheid bar (90). The seed coat remains attached to the funiculus (100). The sub-hilar region contains well-developed vascular tissue (recurrent vascular bundles; 70) and stellate parenchyma (110).

30 At maturity, the seed coat consists of the palisade layer (13), hourglass cells (17), a partially crushed layer of parenchyma (what remains of the outer

-21-

integument), and an endothelium (40). The remnant of the inner integument (30) is often not distinguishable. The tissues of the hilum although compressed, are retained.

- 5 The stages of seed-coat development are also identified in Tables 1 and 2.

Development of the Soybean Seed Cont - Table I

	Epidermis	Hypodermis	Outer Integument	Inner Integument	Endothelium	Status of Embryo
3 daf	simple cuboidal cells		simple parenchyma; no vascularization	2-4 layers	simple cuboidal cells	undifferentiated proembryo
6 daf	not elongated		simple parenchyma; recurrent vascular bundles in subhilar region	5-6 layers; thick walled	simple cuboidal cells	endosperm starting to develop
9 daf	starting to divide and elongate	cells starting to differentiate from parenchyma	upper, thinwalled parenchyma with developing vascular region; lower, thickwalled parenchyma; upper and lower parenchyma show some characteristics of aerenchyma	2-3 cell layers, stretched, starting to compress	simple cuboidal cells	cotyledons starting to develop
12 daf	division and elongation	hourglass cells developing	upper, thinwalled parenchyma with extending vascular region; lower, thickwalled parenchyma, more aerenchyma-like characteristics	1-2 cell layers; more compression	cells starting to stretch	cotyledons shifting
15 daf	cell walls thickening	hourglass cells	upper, thinwalled parenchyma with extending vascular region; lower, thickwalled parenchyma, more aerenchyma-like characteristics	layers of deeply staining wall visible	cells stretching	cotyledons expanding
18 daf	palisade (macroscleireids)	hourglass cells (I-shaped ostioscleireids)	upper, thinwalled parenchyma with extending vascular region; lower, thickwalled parenchyma, more aerenchyma-like characteristics	compressed; one thick, deeply stained line	cells stretching	cotyledons expanding endosperm compressing; protein accumulation starting
21 daf	palisade (macroscleireids)	hourglass cells	upper, thinwalled parenchyma with vascular region completed; lower, thickwalled parenchyma, more aerenchyma-like characteristics	line is thinning	small, oblong, thick-walled cells	endosperm disappearing, protein and lipid accumulation in cotyledons
24 daf	palisade (macroscleireids)	hourglass cells	upper, thinwalled parenchyma with vascular region; lower, thickwalled parenchyma, more aerenchyma-like characteristics	thinning	small, oblong, thick-walled cells	endosperm no longer distinct, protein and lipid accumulation continue
30 daf	palisade (macroscleireids)	hourglass cells	upper and lower parenchyma starting to compress	thin, deeply stained line visible	small, oblong, thick-walled cells	protein and lipid accumulation slowing
45 daf	palisade (macroscleireids)	hourglass cells	parenchyma partially collapsed, upper and lower parenchyma not distinct, vascular tissue not distinguishable	not distinguishable	small, oblong, thick-walled cells	mature cotyledons

SUBSTITUTE SHEET (RULE 26)

Table 2
Position and Levels of Starch, Protein and Lipid in Relation to Seedcoat Development in Soybean

	Developmental Information	Status of Embryo	Starch Accumulation
1 daf	undifferentiated integument and endothelium distinguishable		throughout seedcoat (also in pod, funiculus & senescing floral parts)
3 daf	inner and outer integument, vascular bundles visible		throughout seedcoat (> in outer integument) except epidermis & recurrent vascular bundles (also in funiculus & pod)
6 daf	inner and outer integument, vascular bundles at hilum	embryo starting to expand	outer integument & stellate parenchyma (pod, trichomes & funiculus)
9 daf	inner integument stretched, vascular bundles expanding away from hilum; outer integument differentiating into an upper and lower region	cotyledons are small and beginning to shift	gradient in epidermis; very few granules in bottom half of seedcoat, stellate parenchyma, outer integument
12 daf	inner integument is crushed, epidermis starting to differentiate into palisade and hypodermis starting to differentiate into hourglass cells	cotyledons have shifted, fill embryonic space	palisade (at hilum), stellate parenchyma, differentiating hourglass cells, outer (outer and inner) and inner integument (funiculus) (starting in cotyledons)
15 daf	epidermis differentiated into palisade and hypodermis differentiated into hourglass cells, upper parenchyma differentiated into upper and lower region		palisade, hourglass cells, stellate parenchyma, outer integument (outer region) (cotyledon epidermis)
18 daf	palisade and hourglass cells fully developed, vascular region is very prominent, seed coat is fully expanded	cotyledons have fully expanded	palisade, hourglass, sparse in stellate, sparse in outer integument (outer and inner) (cotyledons)
21 daf	same as in 18		palisade, sparse in stellate and lower portion of upper parenchyma & vascular parenchyma (*also in layer outside of palisade) (cotyledons)
24 daf	same as in 18		palisade, sparse in lower region of upper parenchyma, starting in endothelium (*also in layer outside of palisade) (cotyledons)
30 daf	outer parenchyma region has disappeared, stellate parenchyma present, vascular regions still present, endothelium prominent		endothelium (also endothelium around hypocotyl)
45 daf	outer parenchyma disappeared, stellate parenchyma present, inner integument collapsed, endothelium very prominent		endothelium (also endothelium around hypocotyl)

SUBSTITUTE SHEET (RULE 26)

Table 2 (cont'd)
Position and Levels of Starch, Protein and Lipid in Relation to Seedcoat Development in Soybean

	Protein Accumulation	Lipid accumulation
1 daf	no distinct protein bodies	
3 daf	no distinct protein bodies	few small lipid droplets in funiculus, nothing in seedcoat
6 daf	no distinct protein bodies	few small droplets in outer integument
9 daf	no distinct protein bodies	small droplets in funiculus and counter palisade, sparse in outer integument
12 daf	no distinct protein bodies in seedcoat but starting in cotyledons	
15 daf	protein bodies in cotyledons getting larger	
18 daf		
21 daf		
24 daf		
30 daf	many large and small protein bodies filling cotyledons	
45 daf		

SUBSTITUTE SHEET (RULE 26)

Early development of the tobacco seed coat

At 6 days after anthesis, the tobacco seed coat consists of an epidermis of very large, thin walled cells; a layer of parenchyma cells up to 6 cells thick; and an endothelium of thin-walled, cuboidal cells. By 10 days after anthesis, the inner walls of the epidermis have thickened significantly, with 2-3 layers discernible; the thin-walled parenchyma has become reduced to 3-4 cells thick due to stretching of the layer as the seed expands; and the endothelial cells have become thinner and elongated. At 22 days after anthesis, the epidermal cells have stretched and elongated to accommodate the expanding seed, and the parenchyma and endothelium have elongated and fused into a crushed layer with few individual cells distinguishable.

Seed-coat cryptic promoter

There are several lines of evidence that suggest that the seed-coat expression of GUS activity in the plant T218 is regulated by a cryptic promoter. The region surrounding the promoter and transcriptional start site for the GUS gene are not transcribed in untransformed plants. Transcription was only observed in plant T218 when T-DNA was inserted in *cis*. DNA sequence analysis did not uncover a long open reading frame within the 3.3 kb region cloned. Moreover, the region is very AT rich and predicted to be noncoding (data not shown) by the Fickett algorithm (Fickett, 1982, *Nucleic Acids Res.* **10**, 5303-5318) as implemented in DNASIS 7.0 (Hitachi). Southern blots revealed that the insertion site is within the *N. tomentosiformis* genome and is not conserved among related species as would be expected for a region with an important gene.

As this is the first report of a cryptic promoter specific to seed-coat tissues in plants, it is impossible to estimate the degree to which cryptic

-26-

promoters may contribute to the high frequencies of promoterless marker gene activation in plants. It is interesting to note that transcriptional GUS fusions in *Arabidopsis* occur at much greater frequencies (54%) than translational fusions (1.6%, Kertbundit *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88, 5212-5216).

5 The possibility that cryptic promoters may account for some fusions was recognized by Lindsey *et al.* (1993, *Transgenic Res.* 2, 33-47).

The results disclosed herewith confirms others (Gheysen *et al.*, 1987, *Proc. Natl. Acad. Sci. USA* 84, 6169-6173 and 1991, *Genes Dev.* 5, 287-297) that T-DNA may insert into A-T rich regions as do plant transposable elements (Capel *et al.*, 1993, *Nucleic Acids Res.* 21, 2369-2373). We illustrate that promoters of plant genes are also A-T rich raising speculation that gene insertions into these regions could facilitate the rapid acquisition of new regulatory elements during gene evolution.

15 The insertion of functional genes into the nuclear genome and acquisition of new regulatory sequences has already played a major role in the diversification of certain genes and the endosymbiosis of organelles. In plants, most organellar proteins are nuclear encoded due to the ongoing transfer of their genes into the nucleus (Palmer, 1991, In Bogorad L and Vasil IK (eds) *The Molecular Biology of Plastids*, Academic Press, San Diego, pp 5-53). Recently, it has been shown that the *cox 2* gene of cowpea (Nugent and Palmer, 1991, *Cell* 66, 473-481) and soybean (Covello and Gray, 1992, *EMBO J.* 11, 3815-3820) were transferred from mitochondria to nucleus without promoters by RNA intermediates. The results disclosed herewith, with T-DNA-mediated gene fusions reveal the facility with which promoters can be acquired by incoming genes. The presence of cryptic promoters and diverse regulatory elements in the intergenic regions may insure that genes rapidly achieve the features needed to meet the demands of complex multicellular organisms.

-27-

Other seed-coat, and seed-coat-associated promoters

Transcripts encoding seed coat specific genes were isolated from seed-coat cDNA libraries. These cDNA transcripts were then used to identify the corresponding structural genes and associated promoters from genomic DNA libraries. These promoters, genes and gene products have been isolated and characterized. Examples of such genes include, but are not limited to, SC4, SC20, SC21, a peroxidase cloned from the *Ep* locus, and HP (hydrophobic protein). It is to be understood that this seed-coat library comprises tissues typically found within the seed-coat and tissues adhering to the seed-coat such as the membranous endocarp and cells found in the funicular region such as arial cells (see above for full definition of seed-coat).

Ep locus Peroxidase

The amount of peroxidase activity present in seed coats may vary substantially among different cultivars. The presence of a single dominant gene *Ep* causes a high seed coat peroxidase phenotype. Homozygous recessive *epep* plants are ~100-fold lower in seed coat peroxidase activity which results from a reduction in the amount of peroxidase enzyme present, primarily in the hourglass cells of the subepidermis (Gijzen *et al.*, 1993). In plants carrying the *Ep* gene, peroxidase is heavily concentrated in the hourglass cells (osteosclereids; which form a highly differentiated cell layer with thick, elongated secondary walls and large intercellular spaces).

A seed-coat peroxidase gene, corresponding to the *Ep* locus, was obtained from a soybean seed-coat library. The genomic DNA sequence comprises four exons spanning bp 1533-1752 (exon 1), 2383 -2574 (exon 2), 3605-3769 (exon 3) and 4033-4516 (exon 4) and three introns comprising 1752-2382 (intron 1), 2575-3604 (intron 2) and 3770-4516 (intron 3), of SEQ ID NO:2. Features of the upstream regulatory region of the genomic DNA include a TATA box

-28-

centred on bp 1487; a cap signal 32 bp down stream centred on bp 1520. Also noted within the genomic sequence are three polyadenylation signals centred on bp 4520, 4598, 4663 and a polyadenylation site at bp 4700. The promoter region of the genomic sequence comprises nucleotides 1-1532 of SEQ ID NO:2 (see co-pending US patent application serial No. 08/723,414 and 08/939,905, both of which are incorporated by reference).

Expression of *Ep* is first detected at 6 DPA in the thin-walled parenchyma of the outer integument, adjacent to the thick-walled parenchyma, and flanking the hilum region. By 9 DPA a thin band of expression extends around the entire seed coat, at the junction of the thin-and thick-walled parenchyma. Expression shifts to the hourglass cells as they begin to develop, at 12 DPA (see Figures 13 (e), (f) and (g)).

Expression of a gene under the control of the *Ep* (peroxidase) promoter (nucleotides 1-1532 of SEQ ID NO:2, also see co-pending US patent application serial No. 08/723,414 and 08/939,905, both of which are incorporated by reference) is observed within the seed-coat from 6 to 18 days after anthesis is shown in Figures 13 (d) to (g).

Hydrophobic Protein (HP)

Soybean HP is an 8.3 kD protein consisting of 80 amino acids rich in hydrophobic residues and entirely lacking methionine, phenylalanine, tryptophan, lysine and histidine residues (see Figure 15). The amino acid sequence shows no significant homology to any known proteins (Odani *et al.*, 1987, *Eur J Biochem* 162, 485-491).

To determine the composition of proteins deposited on the soybean seed surface, seeds were washed with a detergent-buffer solution and the extracted peptides were separated by SDS-PAGE. Protein extracts from the seed coat and

-29-

embryo were also prepared for comparison. These results are shown in Figure 17 (a). The embryo and seed coat extracts contained many proteins covering a wide molecular mass range. In contrast, extracts from the seed surface were dominated by a few low molecular mass proteins. Figure 17 (b) demonstrates that HP extraction and separation by SDS-PAGE is dependant on dithiothreitol (DTT).

Even though HP is an abundant seed constituent and a potent allergen, there have been no studies on the expression or localization of the protein or any description of the corresponding gene. This is the first report on the isolation and characterization of HP cDNA (SEQ ID NO:6) and the corresponding genomic clone (SEQ ID NO:7), the pattern of gene expression (Figure 10 (e)), and the localization of the protein (Figure 14 (b)) and its effect on seed luster (Figure 16). Figure 14 (c) shows that the presence of surface protein is related to the luster, or light reflective properties of the seed surface. Surface extracts from shiny seeded phenotypes usually contained far less protein than dull seeded extracts. Moreover, there were large differences in the amount of protein present on the seed surfaces of the two bloom phenotypes examined.

These results also show that the outermost components of the seed coat are in fact derived from the inner layer of the pod wall (see Figure 14 (a)).

The cDNA and genomic copies of the seed-coat associated HP gene were obtained from lambda libraries prepared from cultivar Harosoy 63. The genomic DNA sequence comprises a promoter region from 1-2526 of SEQ ID NO:7. Within this promoter region are located clustered direct repeats (between 1-586; see also Figure 11 (c). and a TATA box located at position 2442-2447. The ORF for HP is between 2526-2882, with the translational start site at 2526, followed by a signal sequence from 2526-2642, and the mature protein from 2643-2882. Also noted within the genomic sequence are six polyadenylation signals and a polyadenylation site at bp 3193.

-30-

Developmental and tissue specific expression patterns for the *HP* gene were determined by RNA blot analysis and *in situ* hybridization. Representative RNA blots, probed with *HP* cDNA, are shown in Figures 10 (e) and (f). These results show that *HP* is highly expressed in the pod during the mid to late stages of seed development. Hybridization signals were also observed in seed coat RNA samples. No expression was evident in the flower, leaf, embryo, stem, or root. We also compared *HP* transcript levels of two different seed luster phenotypes that differed in the amount of HP present on their seed surfaces. Figure 10 (g) shows that *HP* mRNA levels are several fold greater in dull seeded plants that accumulate large amounts of HP on the seed surface when compared to shiny seeded plants that have only trace amount of HP on the seed surface. Faint signals, corresponding to low *HP* transcript levels, were detectable in shiny seeded phenotypes after prolonged exposure times (not shown).

Localization of *HP* gene expression by *in situ* hybridization is shown in Figures 14 (b), (c) and (d). At six days post anthesis (DPA) expression of *HP* is limited to the membranous inner layer of the pericarp. By 12 DPA expression is very strong and the inner epidermis is showing signs of becoming detached from the rest of the pericarp and, in places, is adhering to the seed surface. Tissue sections from this stage of development also showed strong hybridization signals in the sclerenchyma, indicating that *HP* expression occurs throughout the endocarp. Portions of membranous endocarp adhere to the seed during the course of development (see Figures 14 (a) and 16) and thus constitute a newly identified component of the seed coat of mature, fully developed soybeans. The deposition of this material alters the physical properties and the composition of the seed surface, as shown by SDS-PAGE analysis (Figures 17 (a), (b) and (c)) and by scanning electron microscopy (Figure 16). A comparison of dull- and shiny-seeded cultivars reveals that the *HP* gene controls this phenotypic trait in soybeans.

-31-

In Figure 14 (b) can be seen the expression of a gene under the control of the HP promoter, The promoter (nucleotides 1-2526 of SEQ ID NO:7) is active within the membranous endocarp associated with the outer seed-coat.

5 SC 4. 20 and 21

Genes expressing specifically in seed coat tissue were isolated from a seed coat cDNA library obtained from seed coats in later stages of development.

10 SC4

The deduced protein sequence from the SC4 cDNA (Figure 19 (a); SEQ ID NO:3) consists of 289 amino acids and has a molecular mass of 31.9 kDa and a predicted pI of 7.95. Three putative glycosylation sites are present at positions 92, 128 and 269. The putative polypeptide encoded by SC4 exhibits similarity with proteins that comprises a BURP domain (see Figure 19 (b)). The BURP domain is a long carboxyl terminal domain containing a number of highly conserved amino acids (Hattori J. et al., 1998. Mol. Gen. Genet. 259: 424-428). The genomic sequence of *sc4* is provided in SEQ ID NO:9 (also see Restriction map Figure 11 (d)) and comprises a promoter from nucleotides 1-5514 of SEQ ID NO:9.

The expression of a gene under control of the SC4 promoter (nucleotides 1-5514 of SEQ ID NO:9) within soybean seed coat at 3 days after anthesis is shown in Figure 13 (a). The activity of the promoter is localized within the inner integument (arrow). Other areas of brightness in this figure include the recurrent vascular bundles in the funiculus, and the trichomes of the pod (the bright areas are due to the birefringence of crystalline areas in the cell walls, and are also present in the negative control; data not shown).

30

-32-

RNA samples from seed coat, embryo, stem, root, leaf, pod and flower were hybridized with a radiolabelled SC4 cDNA probe (Figure 10 (a)) to determine organ specificity of the expression of SC4. The sc4 transcript was only expressed in the seed coat organ. It was estimated that the size sc4 mRNA was 1.2 kb (data not shown).

Northern blot analysis was carried out to determine the temporal expression pattern of sc4. RNA from seed coat, embryo and pod organs between 6-24 dpa were hybridized with a radiolabelled SC4 cDNA probe. No gene expression was observed in any of the embryo development stages examined (Figure 20 (a)). sc4 expression was apparent in the seed by 6 dpa. After 6 dpa the expression of sc4 in the seed coat increased ~4-fold to its maximum detected level between 9-12 dpa. By 15 dpa sc4 expression had decreased by ~2.5-fold dpa and continued to decline to just detectable levels by 18 dpa (Fig. 3.7). Expression of sc4 could only be detected in the seed coat at 21-24 dpa when the filter was over-exposed. Gene expression of sc4 in the pod was detected from 12-21 dpa only after over-exposure of the filter (data not shown).

To analyse the distribution of sc4 expression with respect to cell differentiation during seed coat development *in situ* hybridization was performed on seed sections from 3-24 dpa seeds. sc4 was expressed throughout the inner integument of the seed coat at 3 dpa (Figures 13 (a) and 21). By 6 dpa the expression pattern of sc4 had changed, and was localized to the outer integument parenchyma but not to the vascular tissue embedded within this layer. sc4 expression in the outer integument was maintained until 18 dpa after which time no further expression was detected (see Table 4 in Examples). In concurrence with northern blot analysis, the *in situ* hybridization results revealed that sc4 expression increased to a maximum between 9-12 dpa and decreased thereafter (Table 4, in Examples). In addition, expression of sc4 was not observed in the embryo of seed at 3-6 dpa.

-33-

Expression of a gene under the control of the SC4 promoter (1-5514 of SEQ ID NO:9) is seen in Figures 13 (a) and 21.

Southern blot analysis was carried out to examine the gene family composition of sc4. Soybean genomic DNA was cleaved with Eco RI, Hind III and Xba I. which do not have recognition sites in the SC4c cDNA sequence. Under conditions of low to high stringency (i.e., from 40-10°C below T_m of the probe) the SC4 cDNA probe hybridized to a single band (Figure 22) and therefore sc4 appears to be a single gene.

Southern blot analysis was also performed to determine the occurrence of sc4 within the following plant species: pea (*Pisum sativum*), canola (*Brassica napus*), oat (*Avena sativa*), onion (*Allium cepa*), pepper (*Capsicum annuum*), Mimosa sp. (*Mimosa pudica*), black spruce (*Picea mariana* (Mill) B.S.P.), birch (*Betula pendula* Roth). The genomic DNA was digested with Eco RI. Under all stringency conditions it was observed that the radiolabelled SC4 cDNA probe hybridized to only soybean genomic DNA (Figure 22 (b)). Further analysis of more related species to soybean need to be carried out.

SC20

The open reading frame of SC20 encodes a putative protein of 770 amino acid residues with a calculated molecular mass of 82.688 kDa and a predicted pI of 6.93. The predicted protein has ten potential N-glycosylation sites (Figure 23 (b)). The hydropathy profile (Figure 23 (c)) of SC20 protein revealed that the first 23 amino acids constitute a hydrophobic region typical of an eukaryotic signal peptide. From northern blot analysis, the SC20 cDNA clone hybridizes to a ~2.5 kb transcript.

The genomic sc20 clone is 7235 bp in length (see Figure 23 (a) for restriction map, and SEQ ID NO:8). Alignment of sc20 genomic and SC20

-34-

cDNA sequences revealed that *sc20* contained eight introns of 94 bp, 101 bp, 185 bp, 80 bp, 154 bp, 112 bp, 110 bp and 93 bp respectively (Figure 23 (a)). A search of the 5' upstream region of *sc20* revealed three potential transcription start sites at positions 1085, 1156 and 2272. The promoter region of *sc20* spans nucleotides 1-2450 of SEQ ID NO:8.

Sequence comparisons (Figure 23 (d)) revealed that the putative polypeptide encoded by SC20 was similar to plant proteins belonging to the Pyrolysins family in the clan of serine proteases known as the subtilases (Barrett A.J. and Rawlings N.D., 1995. Arch. Biochem. Biophys. 318:247-250; Siezen, R. J. and Leunissen, J. A. M. 1997. Protein Sci. 6: 501-523.). The SC20 protein comprises 3 domains: a signal peptide of 23 residues followed by a prosequence of 93 residues and a mature domain of 654 residues. The predicted mature domain of SC20 has a calculated molecular weight of 69.918 kDa and an isoelectric point of 6.34.

Northern blot analysis was carried out to determine specificity of *sc20* expression in various soybean organs i.e., seed coat, embryo, stem, root, leaf, pod and flower (Figure 10 (b)). *sc20* has seed coat-specific expression as its mRNA was detected only in the seed coat organ. The *sc20* transcript was determined to be approximately 2.5 kb (data not shown). Even after prolonged exposure of the filter, no *sc20* transcripts were detected in any of the other plant organs.

Northern blot analysis was performed to determine the temporal gene expression pattern of *sc20* in seed coat, embryo and pod organs of soybean. Total RNA prepared from organs between 6- 24 dpa were probed with a radiolabelled SC20 cDNA probe. *sc20* expression was detected at 9 dpa and rose 1.5 fold to its maximum observed level at 12 dpa (Figure 24). By 18 dpa accumulation of *sc20* mRNA had decreased 4-fold. Prolonged exposure of the filter enabled detection of *sc20* expression at 6 dpa and 21-24 dpa. No gene

-35-

expression was observed at any stage of embryo or pod development examined even after prolonged exposure of the filters. This confirmed that sc20 expression was seed coat-specific.

5 *In situ* hybridization was carried out to analyse the spatial gene expression pattern of sc20 within the seed coat between 3-24 dpa. Seed sections were hybridized with radiolabelled sense and anti-sense SC20 RNA probes. No birefringent cell structures were evident in the seed sections used (Figure 24).

10 Gene expression of sc20 was localized to the thick-walled parenchyma of the outer integument (see Figures 13 (b) and 24). The temporal expression pattern of 9-21 dpa expression with an observed peak at 12 dpa was almost identical to that determined by northern blot analysis (Table 4, in Examples). sc20 transcripts were not detected in the embryo between 3-6 dpa. The *in situ*
15 hybridization results of the seed sections concur with the northern blot analysis that within the seed organ sc20 is expressed only in the seed coat organs.

 Expression of gene under control of the SC20 promoter (1-245 of SEQ ID NO:8) is seen in Figures 13 (b) and 24.

20 Southern blot analysis was performed to ascertain whether sc20 is a single gene or a member of a gene family. Soybean genomic DNA was cleaved with Eco RI, Hind III, Xba I and Eco RV which have three, four, two and one recognition site(s) respectively in the sc20 clone (see Figure 23 (a)). Under
25 conditions of high stringency to detect genes with at least 90% similarity to sc20 the probe hybridized to a single band (Figure. 25 (b)). Under medium stringency conditions to observe genes with 80% similarity to sc20 it was observed that the SC20 probe annealed to 2-3 bands for each digest (Figure 25 (a)). Under
30 conditions of low stringency i.e., 40°C below T_m the SC20 probe hybridized to several more bands from each digest (data not shown). This suggested that sc20

-36-

is a member of a small gene family composed of 2-3 members and that the soybean genome contains several genes which are more distantly related to sc20.

Southern blot analysis was also performed to determine the distribution of sc20 among a number of diverse plant species i.e., pea (*Pisum sativum*), canola (*Brassica napus*), oat (*Avena sativa*), onion (*Allium cepa*), pepper (*Capsicum annuum*), Mimosa sp. (*Mimosa pudica*), black spruce (*Picea mariana* (Mill) B.S.P.), birch (*Betula pendula* Roth). The genomic DNA was restricted with Eco RI. The SC20 cDNA probe hybridized to only the genomic DNA of soybean (Figure 25 (c)) irrespective of stringency conditions utilized. It is possible that the gene may exist in more species more closely related to soybean.

SC21

The expression of a gene under the control of SC21 promoter (see Figure 11 (b)) within seed coat tissues at 15 days after anthesis is shown in Figure 13 (c). Note specific localization of the probe in the thin-walled parenchyma of the outer integument, including the area immediately surrounding the tracheid bar (arrow).

The nucleotide sequences of SC21 (SEQ ID NO:5) and SC17 were identical apart from the position of the poly (A) tail and were just less than 65% similar to a *Cicer arietinum* (chickpea) mRNA for an unknown protein.

The expression of genes under the control of seed-coat promoters of this invention are shown in Figures 10, 13, 14, 21 and 24.

The results of these and other experiments indicating the expression patterns of these genes is summarized in Table 4 within the Examples section.

-37-

The promoters of the present invention can be used to control the expression of any given gene spatially and developmentally within developing seed coats, or seed-coat associated tissues. Some examples of such uses, which are not to be considered limiting, include:

5

1. Modification of storage reserve yields in seed coats, such as starch by the expression of yeast invertase to mobilize the starch, or increasing starch levels by increasing the sink strength by enhancing carbon unloading into seeds, by expressing invertase in specific seed coat tissues, or reduce starch levels by inhibit starch biosynthesis through the expression of the antisense transcript of ADP-glucose pyrophosphorylase.
- 10 2. Modification of seed colour contributed by anthocyanin pigments or condensed tannins in the seed coats by expression of antisense transcripts of the phenylalanine ammonia lyase or chalcone synthase genes.
- 15 3. Modification of fibre content in seed-derived meal by expression of antisense transcripts of the caffeic acid-o-methyl transferase or cinnamoyl alcohol dehydrogenase genes.
- 20 4. Inhibition of seed coat maturation by expression of ribonuclease genes to allow for increased seed size, and to reduce the relative biomass of seed coats, and to aid in dehulling of seeds.
- 25 5. Expression of genes in seed coats coding for insecticidal proteins such as α -amylase inhibitor or protease inhibitor.
- 30 6. Partitioning of seed metabolites such as glucosinolates into seed coats for fungal or insect resistance.

-38-

7. Production of high value proteins in seed coats for use as pharmaceuticals or for use in industrial processes.
8. Control of seed borne diseases by expressing antifungal antiviral, or anti-bacterial proteins within the seed coat

Furthermore, modifications of the nucleotide, or amino acid, sequence of HP, or the preparation of chimeric gene constructs comprising the regulatory region of HP associated with a gene of interest will result in:

- alterations in the textural, visual, chemical or other properties of the seed coat, including the seed surface;
- the production of plants that are less susceptible to seed borne and pod diseases by expressing heterologous proteins in tissues of the ovary wall;
- lessening the health hazard of seed dust exposure by genetic selection or transformation, to produce plants with reduced allergenic protein expression on the seed surface

Thus this invention is directed to such promoter and gene combinations. Further this invention is directed to such promoter and gene combinations in a cloning vector, wherein the gene is under the control of a seed coat specific promoter and is capable of being expressed in a plant cell transformed with the vector. This invention further relates to transformed plant cells and transgenic plants regenerated from such plant cells. The promoter and promoter gene combination of the present invention can be used to transform any plant cell for the production of any transgenic plant. The present invention is not limited to any plant species.

-39-

The following list summarises the nucleotide sequence data in the SEQUENCE LISTING of the present application:

pT218 genomic DNA sequence is found in SEQ ID NO:1;

Ep genomic DNA sequence is listed in SEQ ID NO:2;

5 SC4 cDNA sequence is presented in SEQ ID NO:3;

SC20 cDNA sequence is in SEQ ID NO:4;

SC21 cDNA sequence is presented in SEQ ID NO:5;

HP cDNA is listed in SEQ ID NO:6;

HP genomic DNA sequence is found in SEQ ID NO:7;

10 SC20 genomic DNA sequence is listed in SEQ ID NO:8; and

SC4 genomic DNA sequence is presented in SEQ ID NO:9.

While this invention is described in detail with particular reference to preferred embodiments thereof, said embodiments are offered to illustrate but not
15 limit the invention.

EXAMPLES

Characterization of a Seed Coat-Specific GUS Fusion

20

Transfer of binary constructs to *Agrobacterium* and leaf disc transformation of *Nicotiana tabacum* SR1 were performed as described by Fobert *et al.* (1991, *Plant Mol. Biol.* 17, 837-851). Plant tissue was maintained on 100 μ g/ml kanamycin sulfate (Sigma) throughout *in vitro* culture.

25

Nine-hundred and forty transgenic plants were produced. Several hundred independent transformants were screened for GUS activity in developing seeds using the fluorogenic assay. One of these, T218, was chosen for detailed study because of its unique pattern of GUS expression.

30

-40-

Fluorogenic and histological GUS assays were performed according to Jefferson (*Plant Mol. Biol. Rep.*, 1987, 5, 387-405), as modified by Fobert *et al.* (*Plant Mol. Biol.*, 1991, 17, 837-851). For initial screening, leaves were harvested from *in vitro* grown plantlets. Later flowers corresponding to developmental stages 4 and 5 of Koltunow *et al.* (*Plant Cell*, 1990, 2, 1201-1224) and beige seeds, approximately 12-16 dpa (Chen *et al.*, 1988, *EMBO J.* 7, 297-302), were collected from plants grown in the greenhouse. For detailed, quantitative analysis of GUS activity, leaf, stem and root tissues were collected from kanamycin resistant F1 progeny of the different transgenic lines grown *in vitro*. Floral tissues were harvested at developmental stages 8-10 (Koltunow *et al.*, 1990, *Plant Cell* 2, 1201-1224) from the original transgenic plants. Flowers of these plants were also tagged and developing seeds were collected from capsules at 10 and 20 dpa. In all cases, tissue was weighed, immediately frozen in liquid nitrogen, and stored at -80°C.

Tissues analyzed by histological assay were at the same developmental stages as those listed above. Different hand-cut sections were analyzed for each organ. For each plant, histological assays were performed on at least two different occasions to ensure reproducibility. Except for floral organs, all tissues were assayed in phosphate buffer according to Jefferson (1987, *Plant Mol. Biol. Rep.* 5, 387-405), with 1 mM X-Gluc (Sigma) as substrate. Flowers were assayed in the same buffer containing 20% (v/v) methanol (Kosugi *et al.*, 1990, *Plant Sci.* 70, 133-140).

Tissue-specific patterns of GUS expression were only found in seeds. For instance, GUS activity in plant T218 (Figure 1) was localized in seeds from 9 to 17 days postanthesis (dpa). GUS activity was not detected in seeds at other stages of development or in any other tissue analyzed which included leaf, stem, root, anther, ovary, petal and sepal (Figure 1). Histological staining with X-Gluc revealed that GUS expression in seeds at 14 dpa was localized in seed coats

-41-

but was absent from the embryo, endosperm, vegetative organs and floral organs (results not shown).

5 The seed coat-specificity of GUS expression was confirmed with the more
sensitive fluorogenic assay of seeds derived from reciprocal crosses with
untransformed plants. The seed coat differentiates from maternal tissues called
the integuments which do not participate in double fertilization (Esau, 1977,
Anatomy of Seed Plants. New York: John Wiley and Sons). If GUS activity is
strictly regulated, it must originate from GUS fusions transmitted to seeds
10 maternally and not by pollen. As shown in Table 3, this is indeed the case. As a
control, GUS fusions expressed in embryo and endosperm, which are the
products of double fertilization, should be transmitted through both gametes.
This is illustrated in Table 3 for GUS expression driven by the napin promoter
(BngNAPI, Baszczyński and Fallis, 1990, *Plant Mol. Biol.* 14, 633-635) which is
15 active in both embryo and endosperm (data not shown).

-42-

Table 3. GUS activity in seeds at 14 days post anthesis.

	Cross		GUS Activity
	♀	♂	nmole MU/min/mg Protein
5			
	T218	T218	1.09 ± 0.39
	T218	WT ^a	3.02 ± 0.19
10	WT	T218	0.04 ± 0.005
	WT	WT	0.04 ± 0.005
	NAP-5 ^b	NAP-5	14.6 ± 7.9
	NAP-5	WT	3.42 ± 1.60
15	WT	NAP-5	2.91 ± 1.97

^a WT, untransformed plants

^b Transgenic tobacco plants with the GUS gene fused to the napin, BngNAP1, promoter (Baszczynski and Fallis, 1990, *Plant Mol. Biol.* **14**, 633-635).

Cloning and Analysis of the Seed Coat-Specific GUS Fusion

Genomic DNA was isolated from freeze-dried leaves using the protocol of Sanders *et al.* (1987, *Nucleic Acid Res.* **15**, 1543-1558). Ten micrograms of T218 DNA was digested for several hours with *EcoRI* using the appropriate manufacturer-supplied buffer supplemented with 2.5 mM spermidine. After electrophoresis through a 0.8% TAE agarose gel, the DNA size fraction around 4-6 kb was isolated, purified using the GeneClean kit (BIO 101 Inc., LaJolla,

CA), ligated to phosphatase-treated *EcoRI*-digested Lambda GEM-2 arms (Promega) and packaged *in vitro* as suggested by the supplier. Approximately 125,000 plaques were transferred to nylon filters (Nytran, Schleicher and Schuell) and screened by plaque hybridization (Rutledge *et al.*, 1991, *Mol. Gen. Genet.* 229, 31-40), using the 3' (termination signal) of the *nos* gene as probe (probe #1, Figure 2). This sequence, contained in a 260 bp *SstI/EcoRI* restriction fragment from pPRF-101 (Fobert *et al.*, 1991, *Plant Mol. Biol.* 17, 837-851), was labelled with [α -³²P]-dCTP (NEN) using random priming (Stratagene). After plaque purification, phage DNA was isolated (Sambrook *et al.*, 1989, A Laboratory Manual. New York: Cold Spring Harbor Laboratory Press), mapped and subcloned into pGEM-4Z (Promega). The *EcoRI* fragment and deletions shown in Figure 2 were inserted into pBIN19 (Bevan, 1984, *Nucl. Acid Res.* 12, 8711-8721). Restriction mapping was used to determine the orientation of the fusion in pBIN19 and to confirm plasmid integrity. Plants were transformed with a derivative which contained the 5' end of the GUS gene distal to the left border repeat. This orientation is the same as that of the GUS gene in the binary vector pBI101 (Jefferson, 1987, *Plant Mol. Biol. Rep.* 5, 387-405).

The GUS fusion in plant T218 was isolated as a 4.7 kb *EcoRI* fragment containing the 2.2kb promoterless GUS-*nos* gene at the T-DNA border of pPRF120 and 2.5 kb of 5' flanking tobacco DNA (pT218, Figure 2), using the *nos* 3' fragment as probe (probe #1, Figure 2). To confirm the ability of the flanking DNA to activate the GUS coding region, the entire 4.7 kb fragment was inserted into the binary transformation vector pBIN19 (Bevan, 1984, *Nucl. Acid Res.* 12, 8711-8721), as shown in Figure 2. Several transgenic plants were produced by *Agrobacterium*-mediated transformation of leaf discs. Southern blots indicated that each plant contained 1-4 T-DNA insertions at unique sites. The spatial patterns of GUS activity were identical to that of plant T218. Histologically, GUS staining was restricted to the seed coats of 14 dpa seeds and was absent in embryos and 20 dpa seeds (results not shown). Fluorogenic assays

of GUS activity in developing seeds showed that expression was restricted to seeds between 10 and 17 dpa, reaching a maximum at 12 dpa (Figure 3 (a) and 3 (b)). The 4.7 kb fragment therefore contained all of the elements required for the tissue-specific and developmental regulation of GUS expression.

5

To locate regions within the flanking plant DNA responsible for seed coat-specificity, truncated derivatives of the GUS fusion were generated (Figure 2) and introduced into tobacco plants. Deletion of the region approximately between 2.5 and 1.0 kb, 5' of the insertion site (pT218-2, Figure 2) did not alter expression compared with the entire 4.7 kb GUS fusion (Figures 3b and 4). Further deletion of the DNA, to the *Sna*BI restriction site approximately 0.5 kb, 5' of the insertion site (pT218-3, Figure 2), resulted in the complete loss of GUS activity in developing seeds (Figures 3b and 4). This suggests that the region approximately between 1.0 and 0.5 kb, 5' of the insertion site contains elements essential to gene activation. GUS activity in seeds remained absent with more extensive deletion of plant DNA (pT218-4, Figures 2, 3b and 4) and was not found in other organs including leaf, stem, root, anther, petal, ovary or sepal from plants transformed with any of the vectors (data not shown).

10

15

20

25

30

The transcriptional start site for the GUS gene in plant T218 was determined by RNase protection assays with RNA probe #4 (Figure 2) which spans the T-DNA/plant DNA junction. For RNase protection assays, various restriction fragments from pIS-1, pIS-2 and pT218 were subcloned into the transcription vector pGEM-4Z as shown in Figures 7 and 2, respectively. A 440bp *Hind*III fragment of the tobacco acetohydroxyacid synthase *SURA* gene was used to detect *SURA* and *SURB* mRNA. DNA templates were linearized and transcribed *in vitro* with either T7 or SP6 polymerases to generate strand-specific RNA probes using the Promega transcription kit and [α - 32 P]CTP as labelled nucleotide. RNA probes were further processed as described in Ouellet *et al.* (1992, *Plant J.* 2, 321-330). RNase protection assays were performed as described in Ouellet *et al.*, (1992, *Plant J.* 2, 321-330), using 10-30 μ g of total

-45-

RNA per assay. Probe digestion was done at 30°C for 15 min using 30 µg ml⁻¹ RNase A (Boehringer Mannheim) and 100 units ml⁻¹ RNase T1 (Boehringer Mannheim). Figure 5 shows that two termini were mapped in the plant DNA. The major 5' terminus is situated at an adenine residue, 122 bp upstream of the T-DNA insertion site (Figure 6). The sequence at this transcriptional start site is similar to the consensus sequence for plant genes (C/TTC↓ATCA; Joshi, 1987 *Nucleic Acids Res.* 15, 6643-6653). A TATA box consensus sequence is present 37 bp upstream of this start site (Figure 6). The second, minor terminus mapped 254 bp from the insertion site in an area where no obvious consensus motifs could be identified (Figure 6).

The tobacco DNA upstream of the insertion site is very AT-rich (>75%, see Figure 7). A search for promoter-like motifs and scaffold attachment regions (SAR), which are often associated with promoters (Breyne *et al.*, 1992, *Plant Cell* 4, 463-471; Gasser and Laemmli, 1986, *Cell* 46, 521-530), identified several putative regulatory elements in the first 1.0 kb of tobacco DNA flanking the promoterless GUS gene (data not shown). However, the functional significance of these sequences remains to be determined.

Cloning and Analysis of the Insertion Site from Untransformed Plants

A lambda DASH genomic library was prepared from DNA of untransformed *N. tabacum* SR1 plants by Stratagene for cloning of the insertion site corresponding to the gene fusion in plant T218. The screening of 500,000 plaques with probe #2 (Figure 2) yielded a single lambda clone. The *Eco*RI and *Xba*I fragments were subcloned in pGEM-4Z to generate pIS-1 and pIS-2. Figure 7 shows these two overlapping subclones, pIS-1 (3.0 kb) and pIS-2 (1.1 kb), which contain tobacco DNA spanning the insertion site (marked with a vertical arrow). DNA sequence analysis (using dideoxy nucleotides in both directions) revealed that the clones, pT218 and pIS-1, were identical over a

length of more than 2.5 kb, from the insertion site to their 5' ends, except for a 12 bp filler DNA insert of unknown origin at the T-DNA border (Figure 6 and data not shown). The presence of filler DNA is a common feature of T-DNA/plant DNA junctions (Gheysen *et al.*, 1991, *Gene* 94, 155-163). Gross rearrangements that sometimes accompany T-DNA insertions (Gheysen *et al.*, 1990, *Gene* 94, 155-163; and 1991, *Genes Dev.* 5, 287-297) were not found (Figure 6) and therefore could not account for the promoter activity associated with this region. The region of pIS-1 and pIS-2, 3' of the insertion site is also very AT-rich (Figure 7).

To determine whether there was a gene associated with the pT218 promoter, more than 3.3 kb of sequence contained with pIS-1 and pIS-2 was analyzed for the presence of long open reading frames (ORFs). However, none were detected in this region (data not shown). To determine whether the region surrounding the insertion site was transcribed in untransformed plants, Northern blots were performed with RNA from leaf, stem, root, flower and seeds at 4, 8, 12, 14, 16, 20 and 24 dpa. Total RNA from leaves was isolated as described in Ouellet *et al.*, (1992, *Plant J.* 2, 321-330). To isolate total RNA from developing seeds, 0.5 g of frozen tissue was pulverized by grinding with dry ice using a mortar and pestle. The powder was homogenized in a 50 ml conical tube containing 5 ml of buffer (1 M Tris HCl, pH 9.0, 1% SDS) using a Polytron homogenizer. After two extractions with equal volumes of phenol:chloroform:isoamyl alcohol (25:24:1), nucleic acids were collected by ethanol precipitation and resuspended in water. The RNA was precipitated overnight in 2M LiCl at 0°C, collected by centrifugation, washed in 70% ethanol and resuspended in water. Northern blot hybridization was performed as described in Gottlob-McHugh *et al.* (1992, *Plant Physiol.* 100, 820-825). Probe #3 (Figure 2) which spans the entire region of pT218 5' of the insertion did not detect hybridizing RNA bands (data not shown). To extend the sensitivity of RNA detection and to include the region 3' of the insertion site within the analysis, RNase protection assays were performed with 10 different RNA probes

-47-

that spanned both strands of pIS-1 and pIS-2 (Figure 7). Even after lengthy exposures, protected fragments could not be detected with RNA from 8, 10, 12 dpa seeds or leaves of untransformed plants (see Figure 5 for examples with two of the probes tested). The specific conditions used allowed the resolution of protected RNA fragments as small as 10 bases (data not shown). Failure to detect protected fragments was not due to problems of RNA quality, as control experiments using the same samples detected acetohydroxyacid synthase (*AHAS*) *SURA* and *SURB* mRNA which are expressed at relatively low abundance (data not shown). Conditions used in the present work were estimated to be sensitive enough to detect low-abundance messages representing 0.001-0.01% of total mRNA levels (Ouellet *et al.*, 1992, *Plant J.* 2, 321-330). Therefore, the region flanking the site of T-DNA insertion does not appear to be transcribed in untransformed plants.

Genomic Origins of the Insertion Site

Southern blots were performed to determine if the insertion site is conserved among *Nicotiana* species. Genomic DNA (5 µg) was isolated, digested and separated by agarose gel electrophoresis as described above. After capillary transfer on to nylon filters, DNA was hybridized, and probes were labelled, essentially as described in Rutledge *et al.* (1991, *Mol. Gen. Genet.* 229, 31-40). High-stringency washes were in 0.2 x SSC at 65°C while low-stringency washes were in 2 x SSC at room temperature. In Figure 8, DNA of the allotetraploid species *N. tabacum* and the presumptive progenitor diploid species *N. tomentosiformis* and *N. sylvestris* (Okamuro and Goldberg, 1985, *Mol. Gen. Genet.*, 198, 290-298) were hybridized with probe #2 (Figure 2). Single hybridizing fragments of identical size were detected in *N. tabacum* and *N. tomentosiformis* DNA digested with *HindIII*, *XbaI* and *EcoRI*, but not in *N. sylvestris*. Hybridizations with pIS-2 (Figure 8) which spans the same region but includes DNA 3' of the insertion site yielded the same results. They did not reveal hybridizing bands, even under conditions of reduced stringency, in

-48-

additional *Nicotiana* species including *N. rustica*, *N. glutinosa*, *N. megalosiphon* and *N. debneyi* (data not shown). Probe #3 (Figure 2) revealed the presence of moderately repetitive DNA specific to the *N. tomentosiformis* genome (data not shown). These results suggest that the region flanking the insertion site is unique to the *N. tomentosiformis* genome and is not conserved among related species as might be expected for regions that encode essential genes.

Cloning of seed-coat genes from Soybean:

a) Isolation of seed-coat cDNA clones

A seed coat cDNA library was constructed in Lambda GEM-4 from poly(A)⁺ mRNA isolated from soybean [*Glycine max* (L.) Merrill] seed coats. A sample of the total amplified library was used to sub-clone inserts from the original lambda vector into pBK-CMV (Stratagene). Random clones were selected from this mass excision for plasmid purification and single-run DNA sequencing to construct an expressed sequence tag (EST) database.

For differential screening, an additional cDNA library was constructed from cultivar Maple Presto (*EpEp*) seed coats. The seed coats were harvested from seeds of four fresh weight groups: < 50 mg, 50-100 mg, 150-250 mg and > 250 mg, to represent all developmental stages. Total RNA was isolated from the seed coats using Trizole reagent (BRL) from which poly (A)⁺ RNA was isolated using Oligotex resin (Qiagen). First and second strand cDNAs were synthesized using the Riboclone cDNA synthesis kit and then cloned into a lambda GEM-4 vector (Promega). This seed coat library was differentially screened with positive and negative cDNA probes to identify genes preferentially expressed in the seed coat. The positive probe was derived from poly (A)⁺ mRNA isolated from seed coat tissues while the negative probe was made from poly (A)⁺ mRNA from seedling, flower bud, leaf, pod and root tissue. The cDNA library was screened with cDNA synthesized from RNA using oligo(dT)₁₅

primer, and hybridizations were carried out in Denhardt's solution (Sambrook et al. (1989) Molecular Cloning, Second Edition) at 65°C; wash 4 x 30 minutes 0.1X SSC 0.1% SDS at 65°C.

5 Twenty-one positive clones were identified after plaque purification. The Lambda vector GEM-4 contains a complete pGEM1 plasmid. During the cloning procedure the cDNA is inserted into the Lambda vector at the multicloning site of this plasmid. The entire pGEM1 plasmid, containing the cDNA insert, can be removed from the Lambda vector by digestion with *SpeI* and then can be
10 relegated to form a functional plasmid. Except for SC11 and SC19, the insert was removed from pGEM1 by digestion with *XbaI* and *EcoRI* and ligated into an alternative plasmid vector pGEM4-Z. Following this protocol 21 seed coat clones were used to transform *E. coli* DH5 α . No transformants were obtained with seed coat clones SC7 and SC10 and so these clones were not processed
15 further.

Seed surface proteins were obtained from soybean. A single seed was placed in a 2 mL plastic capped test tube and surface proteins were extracted by adding 0.5 mL of a buffer-detergent solution containing 10 mM Tris-Cl (pH 7.5)
20 0.5% SDS, and 20 mM DTT, and placing the tube in a boiling water bath for 2 min. The contents of the tube were mixed and an aliquot was withdrawn and centrifuged for 5 min at 14,000 g. Freshly prepared loading buffer containing 20 mM DTT was added to the sample and proteins were electrophoretically separated on 15% acrylamide gels in the presence of SDS (see Figure 17) using a
25 modified Laemmli system, as described by Fling and Gregerson (1986, Anal Biochem 155:83-88). Fixation and visualization of the proteins by silver staining followed the method of Blum et al., (1987 Electrophoresis 8: 93-99). The amino terminal of the major peptides (indicated as HPS in Figure 17 (b)) were micro-sequenced from the blotted proteins according to Moos et al., (1988 J. Biol.
30 Chem 263: 6005-6008). The resulting amino acid sequences were identical and matched existing sequences in the GenBank protein database for HP (Odani et

-50-

al., 1987 Eur J. Biochem 162, 485-491). Both peptides had alternative N-terminal residues of Ala or Ile, as has been previously noted for HP. The different electrophoretic mobilities of the two peptides could not be accounted for from the microsequencing analysis, but may be due to differences in glycosylation.

Several different soybean varieties were also compared by SDS-PAGE analysis (see Figure 17 (c)).

To obtain the cDNA transcript of HP, sequences in the seed coat expressed sequence tag database were searched for reading frames corresponding to the HP amino acid sequence. Using this strategy, several identical cDNA transcripts were isolated from the cDNA library obtained from Harosoy 63 seeds described above that included in their reading frames peptide sequences exactly matching HP. The encoded products of these DNA sequences were identified using the BLASTX program at the NCBI site.

b) Characterization of cDNA clones

Sequence analysis

Agarose gel electrophoresis of *Xba*I/*Eco*RI digests of the 19 remaining plasmid clones indicated that the inserts ranged in size from approximately 350bp to 1600bp, including the poly A tail. Inserts of the seed-coat clones were characterized by double stranded dideoxy sequencing of the 5' and 3' ends of the clones. A preliminary classification of the seed coat cDNA clones was made on the basis of sequence homology in the 3' and 5' ends of the clones. Based on sequence similarity with each other these 19 clones were grouped into 7 groups of clones. Sequence similarity was found between four of these groups and GenBank (with proline rich protein and three peroxidase groups). The three remaining groups had no sequence similarity with GenBank. SC4, SEQ ID

-51-

NO:3 (found to be the same as SC1), SC20, SEQ ID NO:4 (found to be the same as SC15), and SC21, SEQ ID NO:5, each represent one clone of each of the three groups which did not exhibit similarity with anything in the GenBank database.

5

SC4

The 1119 bp nucleotide sequence of SC4 (SEQ ID 3, Figure 19 (a); also see Restriction Map Figure 11 (d)) does not represent the full-length cDNA clone as it does not contain an ATG codon for translation initiation. Two typical polyadenylation signals (AATAAA) are located at positions 1096 and 1102. The deduced protein sequence from the SC4 cDNA (Figure 19 (a)) consists of 289 amino acids and has a molecular mass of 31.9 kDa and a predicted pI of 7.95. Three putative glycosylation sites are present at positions 92, 128 and 269.

10

15

The putative polypeptide encoded by SC4 exhibits similarity with proteins that comprises a BURP domain (e.g. RD22, an *Arabidopsis thaliana* dehydration-responsive protein (Yamaguchi-Shinozaki K. and Shinozaki, K. 1993. Mol. Gen. Genet. 238: 17-25); PG1 β , a *Lycopersicon esculentum* polygalacturonase isoenzyme 1 β subunit (Zheng L. et al., 1992. Plant Cell. 4: 1147-1156); Sali3-2, a *Glycine max* L aluminium-induced protein (Ragland M. and Soliman, K.M. 1997. Plant Physiol. 114: 395); USP, a *Vicia faba* unknown seed protein (Baumlein H. et al., 1991. Mol. Gen. Genet. 225: 459-467) and ADR6, a *Glycine max* L auxin-induced protein (Datta N. et al., 1993. Plant Mol. Biol. 21: 859-869); see Figure 19 (b)). The BURP domain is a long carboxyl terminal domain containing a number of highly conserved amino acids (Hattori J. et al., 1998. Mol. Gen. Genet. 259: 424-428). The carboxyl terminal of the conceptual SC4 protein sequence contains the following conserved amino acids which are typical of the BURP domain proteins: two phenylalanine residues, two cysteine residues and four cysteine-histidine motifs which are also in the conserved alignment of CHX10CHX25-27CHX25-26 CH, where X is any amino

20

25

30

-52-

acid (Figure 19 (b)). This BURP domain proteins also share a similar structural make-up of 3-4 domains (Figure 19 (c)) i.e., an amino-terminal domain containing a hydrophobic sequence, a second domain which may or may not be conserved, a third domain consisting of tandem repeats of a short amino acid sequence (not all BURP domain proteins have this domain) and a long carboxyl-terminal BURP domain (Hattori J. et al., 1998. Mol. Gen. Genet. 259: 424-428). The tandem repeats which make up the third domain do not appear to have a common consensus sequence between the different BURP domain proteins. In addition to the BURP domain, the putative SC4 protein shares sequence similarity between its amino terminus and the conserved segment of the second domain possessed by several of the BURP domain proteins (Figure 19 (b)). It was also determined that the SC4 protein has a region containing two copies of the repeated sequence ESRSEXXYAG where X is any amino acid (Figure 19 (a)) which is similar to the structural organization of the third domain of several BURP domain proteins. Due to the extent of structural and sequence similarity between the SC4 protein and the BURP domain proteins it is likely that SC4 also contains a hydrophobic amino terminal if it was full-length.

SC20

The SC20 cDNA clone was sequenced (Figure. 23 (b)) and found to consist of 2447 bp with one 2310 bp open reading frame starting at nucleotide position 13 and ending at 2322. The TAG stop codon may be leaky as plants have tRNAs capable of misreading it. However, any readthrough will be terminated by a second stop codon TGA which is immediately adjacent to UAG. The 3' untranslated region contains one putative polyadenylation signal (AATAAA) located 21 nt after the stop codon.

The open reading frame of SC20 encodes a putative protein of 770 amino acid residues with a calculated molecular mass of 82.688 kDa and a predicted pI of 6.93. The predicted protein has ten potential N-glycosylation sites (Figure 23

-53-

(b)). The hydropathy profile (Figure 23 (c)) of SC20 protein revealed that the first 23 amino acids constitute a hydrophobic region typical of an eukaryotic signal peptide. From northern blot analysis, the SC20 cDNA clone hybridizes to a ~2.5 kb transcript (data not shown). SC20 was used to obtain the genomic clone which was from a soybean cv. Harovinton genomic library.

Sequence comparisons (Figure 23 (d)) revealed that the putative polypeptide encoded by SC20 was similar to a *Picea abies* (black spruce) AF70 protein (Sabala et al., 1997. *Physiol. Plant.* 99: 316-322); cucumisin, (Yamagata Y. et al. 1994 *J. Biol. Chem.* 269: 32725-32731) from *Cucumis melo* L. (musk melon); a pathogen-induced protein, P69B, (Tornerio P. et al., 1997 *J. Biol. Chem.* 272: 14412-12219) from *Lycopersicon esculentum* (tomato) and a nodule-specific protein, Ag12, (Ribeiro A. et al., 1995 *Plant Cell.* 7: 785-794) from *Alnus glutinosa* (black alder). These plant proteins belong to the Pyrolysins family in the clan of serine proteases known as the subtilases (Barrett A.J. and Rawlings N.D., 1995. *Arch. Biochem. Biophys.* 318:247-250; Siezen, R. J. and Leunissen, J. A. M. 1997. *Protein Sci.* 6: 501-523.). The SC20 protein contains the conserved catalytic residues aspartate, histidine and serine as well as the highly conserved asparagine residue which is involved in stabilizing substrate binding. Moreover, the order of these four conserved residues in the SC20 protein is also a characteristic feature of subtilases. The SC20 protein also has a large sequence insertion between the conserved asparagine and serine residues found in plant subtilases but not in other subtilase members such as subtilisin BPN' (Power S.D. et al., 1986 *PNAS.* 83:3096-3100), kex2 (Mizuno K. et al., 1988 *Biochem. Biophys. Res. Comm.* 156: 246-254) or furin (van de Ven W.J.M. et al., 1990 *Mol. Biol. Rep.* 14: 265-275). Based on sequence similarity between the N-termini of mature plant subtilases and the SC20 protein (data not shown) it was predicted that the SC20 protein had a mature domain starting from residue 117. Therefore the SC20 protein appears to be composed of 3 domains: a signal peptide of 23 residues followed by a prosequence of 93 residues and a

-54-

mature domain of 654 residues. The predicted mature domain of SC20 has a calculated molecular weight of 69.918 kDa and an isoelectric point of 6.34.

HP

The cDNA sequence for HP (pHPScDNA1) is (SEQ ID NO:6) shown in Figure 15. The 700 bp transcript includes 30 bp of 5' untranslated region (UTR), an open reading frame (ORF) of 119 amino acids, and 313 bp of 3' UTR. Several polyadenylation signals were identified in the 3' UTR. The final 80 residues of deduced amino acid sequence exactly match the peptide sequence reported for the hydrophobic protein (Odani *et al.*, 1987, *Eur J Biochem* 162, 485-491). Thus, the HP cDNA transcript indicates that hydrophobic protein is translated with a leader sequence of 39 amino acids.

Northern blot analysis

Northern analysis, using the cDNA inserts of each clone as a probe, was performed to investigate the expression pattern of the 19 seed coat clones.

RNA isolation from leaf, stem, pod and flower tissue was optimized based on a protocol adapted from Tripure Isolation reagent kit (Boehringer Mannheim). Plant tissue was frozen in liquid nitrogen and homogenized with the Tripure reagent (a monophasic solution of phenol and guanidine thiocyanate). After the addition of chloroform the sample is centrifuged so that it separates into three phases. RNA is recovered from the upper aqueous phase by isopropanol precipitation. Due to the problem of polysaccharide contamination which increases with seed maturity, the isopropanol precipitation step was carried out in the presence of high salt which effectively maintains the polysaccharides in a soluble form whilst the RNA is precipitated.

-55-

Total RNA from seed-coat, embryo and root tissue was isolated as described by Fobert et al. (Plant J. 1994 6:567-577). Plant tissue was frozen in liquid nitrogen and homogenized in 1M Tris-HCl, pH9, 1% SDS buffer. The sample was extracted twice with equal volume phenol:chloroform:isoamyl alcohol (25:24:1), nucleic acids were collected by ethanol precipitation, collected by ethanol precipitation and resuspended in water. The RNA was precipitated overnight in 2M LiCl at 0°C, collected by centrifugation and resuspended in water.

RNA was denatured and size fractionated by formaldehyde gel electrophoresis and transferred onto nylon filters. Northern hybridization was carried out using radioactively labeled cDNA probes with hybridization in modified Church's buffer (Church and Gilbert (1994) PNAS USA 81: 1991-1995) at 65°C; wash 2 x 30 minutes 0.1X SSC 0.1% SDS at 65°C. From this analysis, it was observed that SC4, and SC20 have seed coat specific expression. *Ep* locus peroxidase has preferential expression within seed-coat tissues, and SC21 was only expressed in seed coat, stem, root and flower tissues. The results are shown in Figure 10 (a) - (d).

HP

For the analysis of HP expression, total RNA was isolated from roots, stems, leaves, flowers, pods, seed coats, and embryos dissected from soybean plants at various stages of development, according to published methods (Wang and Vodkin (1994) *Plant Mol Biol Rep* 12, 132-145). The RNA samples were quantitated by measuring absorbance at 260 nm, and by electrophoretic separation in formaldehyde gels and comparison to known standards. Samples of total RNA (10 µg each) were blotted to nylon membrane using a vacuum manifold apparatus and fixed by UV cross-linking. Digoxigenin-labelled cDNA was prepared according to instructions of the manufacturer (Boehringer) and used to probe the RNA blots. Results, Figures 10 (e) and (f) show that the *HP* gene is

-56-

highly expressed in the pod tissues during the later stages of development. Hybridization signals were also noted in RNA samples derived from seed coat tissue, but not in RNA samples from the leaf, flower, embryo, stem or root. These results, together with the data from the *in situ* hybridizations (see below) and the scanning electron microscopy analysis, indicate the *HP* gene is specifically expressed in the endocarp of the ovary wall. Pieces of this tissue detach from the pod wall and adhere to the seed surface during development, thus becoming a component of the mature seed coat (see Figure 14 (a)).

SC4

RNA samples from seed coat, embryo, stem, root, leaf, pod and flower were hybridized with a radiolabelled SC4 cDNA probe (Figure 10 (a)) to determine organ specificity of the expression of SC4. The *sc4* transcript was only expressed in the seed coat organ. It was estimated that the size *sc4* mRNA was 1.2 kb (data not shown).

Northern blot analysis was carried out to determine the temporal expression pattern of *sc4*. RNA from seed coat, embryo and pod organs between 6-24 dpa were hybridized with a radiolabelled SC4 cDNA probe. At 6 dpa the seed is too small to separate the seed coat and embryo organs and so total RNA was isolated from an entire seed. *sc4* expression was already apparent in the seed by 6 dpa. No gene expression was observed in any of the embryo development stages examined (Figure 20 (a)). *sc4* mRNA transcripts were not observed in the embryo of 3-6 dpa seed sections examined by *in situ* hybridization using a radiolabelled SC4 antisense RNA probe (data not shown). Therefore the *sc4* expression observed at 6 dpa in the seed tissue is likely to be seed coat derived. After 6 dpa the expression of *sc4* in the seed coat increased ~4-fold to its maximum detected level between 9-12 dpa. By 15 dpa *sc4* expression had decreased by ~2.5-fold dpa and continued to decline to just detectable levels by 18 dpa (Fig. 3.7). Expression of *sc4* could only be detected

-57-

in the seed coat at 21-24 dpa when the filter was over-exposed. Gene expression of sc4 in the pod was detected from 12-21 dpa only after over-exposure of the filter (data not shown).

5 SC20

10 Northern blot analysis was carried out to determine specificity of sc20 expression in various soybean organs i.e., seed coat, embryo, stem, root, leaf, pod and flower (Figure 10 (b)). sc20 has seed coat-specific expression as its mRNA was detected only in the seed coat organ. The sc20 transcript was determined to be approximately 2.5 kb (data not shown). Even after prolonged exposure of the filter, no sc20 transcripts was detected in any of the other plant organs.

15 Northern blot analysis was performed to determine the temporal gene expression pattern of sc20 in seed coat, embryo and pod organs of soybean. Total RNA prepared from organs between 6- 24 dpa were probed with a radiolabelled SC20 cDNA probe. sc20 expression was detected at 9 dpa and rose 1.5 fold to its maximum observed level at 12 dpa (Figure 24). By 18 dpa
20 accumulation of sc20 mRNA had decreased 4-fold. Prolonged exposure of the filter enabled detection of sc20 expression at 6 dpa and 21-24 dpa. No gene expression was observed at any stage of embryo or pod development examined even after prolonged exposure of the filters. This confirmed that sc20 expression was seed coat-specific.

25

b) In situ hybridization

30 To analyze the distribution of the clones mRNA expression with respect to cell differentiation during development, *in situ* hybridization, on sections from 3, 6, 9, 12, 15, 18, 21 and 24 DAF seeds was used. Seeds or parts of seeds were fixed in FAA fixative (50% ethanol, 5% acetic acid and 3.7% formaldehyde),

-58-

dehydrated in an ethanol/ tertiary butyl alcohol series and infiltrated and embedded in paraplast plus. Sections (5-10 μ m) were cut on a microtome, transferred onto Superfrost slides which are positively charged to allow better adherence of the sections to the slide surface. Prior to *in situ* hybridization the samples were dewaxed in a xylene/ethanol series. *In situ* hybridization was carried out with ³⁵S-labelled cDNA sense and anti-sense probes following the method of Cox and Goldberg (1998).

Ep

For the *in situ* analysis of *Ep* expression, flowers were tagged on days of full anthesis when the banner petal was fully extended and harvested at three day intervals from 1-30 days post anthesis (DPA), and at 45 DPA [19]. Tissue samples were fixed in a solution of 3.7% formaldehyde, 50% ethanol, and 5% acetic acid for 3 h at room temperature, dehydrated in an ethanol series (50, 60, 70, 80, 90, 95, 100%) then infiltrated with t-butyl alcohol (TBA) in ethanol in a stepwise series (25, 50, 75, and 100%), followed by infiltration with Paraplast and several changes of pure melted Paraplast at 57 °C. After infiltration, samples were placed in blocks and allowed to harden. Sections of 8-10 μ m were cut on a rotary microtome and affixed to glass slides. Prior to hybridization, sections were de-waxed in xylene, and re-hydrated in an ethanol series (100, 95, 85, 70, 50, 30, 15, 0% ethanol in distilled RNase free water).

Localization of RNA was performed with ³⁵S-labelled RNA probes generated from *Ep* cDNA clones. The prehybridization, hybridization, and wash conditions followed published methods (Cox K.H., and Goldberg R.B. 1988, Analysis of plant gene expression. In Shaw CH (ed), Plant Molecular Biology: A Practical Approach, pp. 1-35. IRL Press, Oxford). Briefly, sections were treated with Proteinase K and acetylated with acetic anhydride in triethanolamine. The sections were then hybridized with ³⁵S-RNA probes overnight at 42 °C, washed, and dehydrated in an ethanol series before application of Kodak NTB-2

-59-

track emulsion. After 1 week at 4 °C, slides were developed in Kodak D-19 developer, fixed in Kodak Fix, and stained in Toluidine Blue O (0.05 % in 50 mM Acetate buffer, pH 4.5). Slides were then dehydrated in an ethanol and xylene series, and mounted in Permount. Slides were photographed on Kodak EPL 400 slide film, using dark field optics

The expression of a gene under the control of the *Ep* (peroxidase) promoter (nucleotides 1-1532 of SEQ ID NO:2, also see co-pending US patent application serial No. 08/723,414 and 08/939,905, both of which are incorporated by reference) is localized within the hourglass cells (arrow; Figure 13(d)) within the seed-coat at 18 days after anthesis. Expression of *Ep* is first detected at 6 DPA in the thin-walled parenchyma of the outer integument, adjacent to the thick-walled parenchyma, and flanking the hilum region (Figure 13 (e)). By 9 DPA a thin band of expression extends around the entire seed coat, at the junction of the thin-and thick-walled parenchyma (Figure 13 (f)). Expression shifts to the hourglass cells as they begin to develop, at 12 DPA (Figure 13 (g)).

HP

For the analysis of HP (Figures 14 (c) and (d)), tissue samples were fixed in a solution of 50 % ethanol, 5 % acetic acid, 3.7 % formaldehyde for 3 h at room temperature, dehydrated in an ethanol series (50, 60, 70, 80, 90, 95, 100 %) then infiltrated with t-butyl alcohol (TBA) in a stepwise series (25, 50, 75, and 100 % TBA in ethanol), followed by infiltration with Paraplast by gradual addition of increasing amounts of Paraplast to 100 % TBA, followed by several changes of pure melted Paraplast at 57 °C. After infiltration, samples were placed in blocks and allowed to harden. Sections of 8 to 10 μ m were cut on a rotary microtome and affixed to glass slides. Prior to hybridization, sections were de-waxed in xylene, and re-hydrated in an ethanol series (100, 95, 85, 70, 50, 30, 15, 0 % ethanol in distilled RNase free water). Sections were then treated with

-60-

Proteinase K and acetylated with acetic anhydride in triethanolamine. Sections were hybridized with ³⁵S-RNA probes overnight at 42 °C, then washed and dehydrated in an ethanol series before application of Kodak NTB-2 track emulsion. After 1 week at 4 °C, slides were developed in Kodak D-19 developer, fixed in Kodak Fix, and briefly stained in Toluidine Blue O before dehydrating in an ethanol and xylene series, then mounting in Permout. Slides were photographed on Kodak EPL 400 slide film, using dark field optics.

The expression of a gene under the control of the HP promoter (nucleotides 1-2526 of SEQ ID NO:7) is localized within the membranous endocarp (arrow, Figure 14 (b)) at 12 days after anthesis. At six days post anthesis (DPA) expression of *HPS* is limited to the membranous inner layer of the pericarp. By 12 DPA expression is very strong and the inner epidermis is showing signs of becoming detached from the rest of the pericarp and, in places, is adhering to the seed surface. Tissue sections from this stage of development also showed strong hybridization signals in the sclerenchyma, indicating that *HP* expression occurs throughout the endocarp.

SC4

To analyse the distribution of *sc4* expression with respect to cell differentiation during seed coat development *in situ* hybridization was performed on seed sections from 3-24 dpa seeds. The seed sections were hybridized with radiolabelled sense and antisense SC4 RNA probes which were detected by exposure of the sections to photographic emulsion. Within the seed sections the antisense or sense RNA probes can be localized by observing the accumulation of silver grains (produced in the emulsion by the emitted β -particles) under dark-field illumination with a light microscope. Cell walls of some plant structures can be birefringent (i.e., reflect light) under dark-field illumination. Two birefringent areas can be observed in both the hilum and the funiculus of the seed sections in

-61-

Figure 21 therefore any expression or lack there-of by sc4 will be masked in these locations.

sc4 was expressed throughout the inner integument of the seed coat at 3 dpa (Figure 21). By 6 dpa the expression pattern of sc4 had changed, and was localized to the outer integument parenchyma but not to the vascular tissue embedded within this layer. sc4 expression in the outer integument was maintained until 18 dpa after which time no further expression was detected (see Table 4 below). In concurrence with northern blot analysis, the *in situ* hybridization results revealed that sc4 expression increased to a maximum between 9-12 dpa and decreased thereafter (Table 4). In addition, expression of sc4 was not observed in the embryo of seed at 3-6 dpa.

The expression of a gene under control of the SC4 promoter (nucleotides 1-5514 SEQ ID NO:9) within soybean seed coat at 3 days after anthesis is also shown in Figure 13 (a). Expression is localized within the inner integument (arrow; Figure 13 (a)). Other areas of brightness in this figure include the recurrent vascular bundles in the funiculus, and the trichomes of the pod (the bright areas are due to the birefringence of crystalline areas in the cell walls, and are also present in the negative control; data not shown).

SC20

In situ hybridization was carried out to analyse the spatial gene expression pattern of sc20 within the seed coat between 3-24 dpa. Seed sections were hybridized with radiolabelled sense and anti-sense SC20 RNA probes. No birefringent cell structures were evident in the seed sections used (Figure 24).

Gene expression of sc20 was localized to the thick-walled parenchyma of the outer integument (see Figures 13 (b) and 24). The temporal expression pattern of 9-21 dpa expression with an observed peak at 12 dpa was almost

-62-

identical to that determined by northern blot analysis (Table 4, in Examples).
sc20 transcripts were not detected in the embryo between 3-6 dpa. The *in situ*
hybridization results of the seed sections concur with the northern blot analysis
that within the seed organ sc20 is expressed only in the seed coat organs.

5

Expression of gene under control of the SC20 promoter (1-2450 of SEQ
ID NO:8) is seen in Figures 13 (b) and 24.

SC21

10

The expression of a gene under the control of SC21 (see Figure 11 (b))
within seed coat tissues at 15 days after anthesis is localized in the thin-walled
parenchyma of the outer integument, including the area immediately surrounding
the tracheid bar (arrow; Figure 13 (c)).

15

c) Construction of genomic libraries

In order to characterise the gene corresponding to seed coat cDNA
clone(s), several genomic libraries were constructed in λ vectors from total DNA
isolated from etiolated seedlings of various soybean cultivars. Two soybean
genomic libraries were constructed in λ Lambda FixII (Stratagene, La Jolla, CA)
from the total DNA isolated from etiolated seedlings of soybean [*Glycine max*
(L.) Merrill] cvs. Harosoy 63 and Harovinton. The DNA was partially digested
with Bgl II prior to ligation into the cloning vector.

25

Genomic clones corresponding to the cDNA clone SC4 and SC20 were
obtained. Lambda DNA was isolated from each plaque. An ~ 8 kb Xba I
fragment from the SC20 lambda clone and an ~ 8 kb Sac I fragment from the
SC4 lambda clone, identified by southern blotting, were ligated into pBlueScript-
SK (Stratagene, La Jolla, CA) and transformed into *E. coli* TOP 10 cells.

30

-63-

Southern blot analysis of genomic soybean DNA, was carried out with 7 seed coat cDNA probes to determine similarities between clones and whether the clones represent a single gene or a gene family. Southern blots were also performed to determine the occurrence of the seed-coat specific genes within other dicotyledonous and monocotyledonous plant species. Soybean genomic DNA was cleaved with several restriction enzymes and the resulting DNA fragments were size fractionated using agarose gel electrophoresis, denatured and transferred to nylon filters. Hybridization was carried out with radiolabelled cDNA probes.

Isolation of genomic clones

Initially, soybean genomic libraries were screened for the presence of the seed coat clone using the polymerase chain reaction with primers specifically designed from each cDNA sequence. This helped to target potential libraries for the isolation of genomic clones. The chosen genomic library was then screened using nucleic acid hybridization with cDNA probes. For genomic library screening hybridization conditions involved using modified Church's buffer (Church and Gilbert (1994) PNAS USA 81: 1991-1995) at 65°C; wash 0.1X SSC 0.1% SDS at 52-55°C. Probes were random primed in presence of ³²PdCTP using standard protocols.

Ep

A seed-coat peroxidase gene, corresponding to the *Ep* locus, was obtained from a soybean seed-coat library. The genomic DNA sequence comprises four exons spanning bp 1533-1752 (exon 1), 2383 -2574 (exon 2), 3605-3769 (exon 3) and 4033-4516 (exon 4) and three introns comprising 1752-2382 (intron 1), 2575-3604 (intron 2) and 3770-4516 (intron 3), of SEQ ID NO:2. Features of the upstream regulatory region of the genomic DNA include a TATA box centred on bp 1487; a cap signal 32 bp down stream centred on bp 1520. Also

-64-

noted within the genomic sequence are three polyadenylation signals centred on bp 4520, 4598, 4663 and a polyadenylation site at bp 4700. The promoter region of the genomic sequence comprises nucleotides 1-1532 of SEQ ID NO:2 (see co-pending US patent application serial No. 08/723,414 and 08/939,905, both of which are incorporated by reference).

HP

For the isolation of the genomic *HP* gene, a genomic library was constructed from DNA isolated from the soybean cultivar Harosoy 63. The DNA was purified and partially digested with *Bgl* II prior to ligation into the cloning vector lambda FixII (Stratagene). The resulting library was amplified and screened with the hydrophobic protein cDNA probe (pHPScDNA1). A positive clone was identified, purified, and found to contain a 14 kb insert. The entire insert was sub-cloned into pBluescript KS(+) and named pHPS1. The *HP* gene was determined by PCR analysis to lie near one end of the 14 kb *Bgl* II fragment (for restriction map see Figure 11 (c)). This region of the pHPS1 insert was sequenced by primer walking, and 3368 bp of this sequence data is disclosed here (SEQ ID NO:7). Aside from the polyadenylation site, the cDNA sequence (pHPScDNA1) exactly matches a stretch of sequence encoded on the genomic clone (pHPS1), indicating that this gene contains no introns. Additionally, a TATA box consensus signal was identified 81 bp upstream from the ATG translation start site.

SC4

A genomic clone corresponding to SC4 cDNA clone was isolated from the soybean genomic library Harosoy 63 (*Bgl* II digest). The genomic sc4 clone is 8310 bp in length (SEQ ID NO:9). The promoter region is found between nucleotides 1-5514 of SEQ ID NO:9. The restriction map is provided in Figure 11 (d).

SC20

A genomic clone corresponding to SC20 cDNA clone was isolated from soybean genomic library prepared from cv Harovinton (GigapackGold packaging). The genomic sc20 clone is 7235 bp in length (see Figure 23 (a), SEQ ID NO:8). Alignment of sc20 genomic and SC20:2 cDNA sequences revealed that sc20 contained eight introns of 94 bp, 101 bp, 185 bp, 80 bp, 154 bp, 112 bp, 110 bp and 93 bp respectively (Figure 23 (a)). A search (www.hgc.lbl.gov/cgi-bin/promoter.pl) of the 5' upstream region of sc20 revealed three potential transcription start sites at positions 1085, 1156 and 2272. The promoter region is found between nucleotides 1-2450 of SEQ ID NO:8. The restriction map of SC20 is presented in Figure 11 (a) and 23(a).

SC21

A genomic clone corresponding to SC21 cDNA clone was isolated from the soybean genomic library prepared from Harosoy 63 (*Eco*R1 digest). The DNA of the SC21 genomic clone was digested with several restriction enzymes, fractionated by agarose gel electrophoresis and transferred onto nylon membrane. Hybridizations were carried out using radiolabelled cDNA. A restriction map of this clone is presented in Figure 11 (b).

Southern analysis

SC4

Southern blot analysis was carried out to examine the gene family composition of sc4. Soybean genomic DNA was cleaved with *Eco* RI, *Hind* III and *Xba* I. which do not have recognition sites in the SC4c cDNA sequence. Under conditions of low to high stringency (i.e., from 40-10°C below *T_m* of the

-66-

probe) the SC4 cDNA probe hybridized to a single band (Figure 22) and therefore sc4 appears to be a single gene.

SC20

Southern blot analysis was performed to ascertain whether sc20 is a single gene or a member of a gene family. Soybean genomic DNA was cleaved with Eco RI, Hind III, Xba I and Eco RV which have three, four, two and one recognition site(s) respectively in the sc20 clone (see Figure 23 (a)).

Hybridization was carried out with radiolabelled SC20 cDNA probe which could anneal from the middle of exon 6 to the Eco RI site on exon 9. For each digest the probe was expected bind to only one of the resulting sc20 restriction fragments. Under conditions of high stringency to detect genes with at least 90% similarity to sc20 the probe hybridized to a single band (Figure. 25 (b)). Under medium stringency conditions to observe genes with 80% similarity to sc20 it was observed that the SC20 probe annealed to 2-3 bands for each digest (Figure 25 (a)). Under conditions of low stringency i.e., 40°C below T_m the SC20 probe hybridized to several more bands from each digest (data not shown). This suggested that sc20 is a member of a small gene family composed of 2-3 members and that the soybean genome contains several genes which are more distantly related to sc20.

Southern blot analysis was performed to determine the occurrence of the seed-coat genes within the following plant species: pea (*Pisum sativum*), canola (*Brassica napus*), oat (*Avena sativa*), onion (*Allium cepa*), pepper (*Capsicum annuum*), mimosa (*Mimosa pudica*), black spruce (*Picea mariana* (Mill B.S.P.)), birch (*Betula pendula* Roth). Genomic DNA was cleaved with *EcoRI* and the resulting DNA fragments were fractionated using agarose gel electrophoresis, denatured and transferred to nylon filters. Hybridization was carried out with radiolabelled SC4 (Figure 22 (b)), SC20 (Figure 25 (c)), SC21, *Ep* locus peroxidase, and HP cDNA probes, using modified Church's buffer at 65°C. The

-67-

filters were washed with 2XSSC, 0.1%SDS at 42°C for 30 minutes, followed by 0.1XSSC, 0.1%SDS at 65°C for 30 minutes. SC4, SC20 and *Ep* locus peroxidase cDNA hybridized to the genomic DNA of soybean only. SC21 cDNA hybridized to the genomic DNA of both soybean and oat. HP cDNA hybridized to the genomic DNA of soybean.

Analysis of promoter activity

The developmental expression of genes under the control of SC4, SC20 SC21 and the peroxidase promoter were further characterized during development of the seed coat by *in situ* hybridization as described above. The results are summarized in Table 4.

Developmental analysis of SC20 indicates that the promoter is highly active at 12 DAF within the outer integument and thick walled parenchyma, however, activity of the SC20 promoter is detectable from about 9 DAF (as per Figure 13 (b)) to about 18 DAF, and is partially detected at 21 DAF.

The SC4 promoter is active from about 3 daf (also see Figure 13 (a)) to about 6 DAF within the inner integument, and then is highly active at 9 DAF within the outer integument and stellate parenchyma, and strongly active at 12 DAF in these same tissues. The SC4 promoter is still active within the outer integument up to 18 DAF.

The SC21 promoter is active throughout seed coat development during all stages examined, from 3 about DAF to about 24 DAF, with strongest activity noted from about 9 DAF to about 15 DAF (also see Figure 14 (c)). The gene under the control of the SC21 promoter is expressed primarily within the outer integument and derived tissues.

-68-

5 The Ep (peroxidase, see co-pending US patent application serial No. 08/723,414 and 08/939,905, both of which are incorporated by reference) promoter is active from about 6 DAF to about 24 DAF. Expression of the peroxidase gene, from about 12 DAF to about 24 DAF, is predominantly within cells of the outer integument, and the hourglass cells (see also Figure 13 (d)).

10 The HP promoter is active from about 9 daf through to about 24 daf. The promoter is active within the membranous endocarp throughout this period of time (see also Figure 14 (b)).

-69-

Table 4
Radioactive *in situ* Hybridization (³²S) of Soybean Seed Coat Tissue (*Glycine max* var. Maple Presto):
Developmental study with seed coat specific clones and peroxidase clones

	SC 20	SC 4	SC 21	Ep	IIP
3 daf	-	++ (inner integument)	+ (outer integument, subhilum region)	-	-
6 daf	-	++ (outer integument)	+(thin-walled outer integument except vascular layer; gradient from hilum to bottom of seed)	+ (localized beneath recurrent vascular bundles)	-
9 daf	+ (outer integument, thick walled parenchyma)	+++ (outer integument except vascular layer)	++ (thin-walled outer integument except vascular layer; gradient from hilum to bottom of seed)	+ (outer integument; thin walled parenchyma beneath vascular tissue)	++ membranous endocarp of the pod
12 daf	++ (outer integument, thick walled parenchyma)	+++ (outer integument except vascular layer)	++ (thin-walled outer integument except vascular layer)	++ (outer integument except vascular layer; hourglass cells)	++ membranous endocarp of the pod
15 daf	+ (outer integument; thick & thin walled parenchyma except vascular layer)	+ (outer integument except vascular layer)	++ (thin-walled outer integument except vascular layer)	++ (outer integument except vascular layer; hourglass cells)	++ membranous endocarp of the pod
18 daf	+ (outer integument; thick & thin walled parenchyma except vascular layer)	(+)	++ (thick & thin walled parenchyma of outer integument except vascular layer)	++ (outer integument except vascular layer; hourglass cells)	++ membranous endocarp of the pod
21 daf	(+)	-	++ (thick & thin walled parenchyma of outer integument except vascular layer)	++ (outer integument except vascular layer; hourglass cells)	++ membranous endocarp of the pod
24 daf	-	-	++ (thick & thin walled parenchyma of outer integument except vascular layer)	++ (outer integument except vascular layer; hourglass cells)	++ membranous endocarp of the pod

+++ highly expressed, many silver grains; ++ moderate expression; + low expression; (+) fading expression; - no expression distinguishable

SUBSTITUTE SHEET (RULE 26)

WEST

Seed Surface Analysis of Dull and Shiny Soybean Varieties

Seed surface proteins of several different soybean varieties were compared by SDS-PAGE analysis. A single seed was placed in a 2 mL plastic capped test tube and surface proteins were extracted by adding 0.5 mL of a buffer-detergent solution (10 mM Tris-Cl (pH 7.5) 0.5% SDS, 20 mM DTT) and placing the tube in a boiling water bath for 2 min. The contents of the tube were mixed and a sample was withdrawn and centrifuged for 5 min at 14,000 g. The proteins in the supernatant were electrophoretically separated on 15% acrylamide gels in the presence of SDS (Fling and Gregerson (1986) *Anal Biochem* 155, 83-88) and detected by silver staining. This analysis revealed that the 8.3 kD hydrophobic protein is by far the most abundant protein molecule occurring on the seed surface of 'Dull' seeded varieties. Only trace amounts of hydrophobic protein was detected on the surface of 'Shiny' seeded soybean varieties (results not shown).

Analysis of seed coat tissues using light microscopy indicated that the membranous endocarp of the pod wall remains in association with the seed-coat (Figure 14 (a)). Scanning electron microscopy (SEM) of the seed surface of soybeans also showed obvious differences between 'Dull' (e.g. cultivar Clark) and 'Shiny' (e.g. cultivar Williams 82) varieties (see Figure 16). Whole seeds were sputter coated with gold and examined by SEM at several magnifications. When viewed with the naked eye, 'Dull' varieties present a surface with a powder-like coating whereas 'Shiny' types appear to have a smoother and more light-reflective surface. Examination by SEM at low magnification (18 X) reveals that the surface of 'Dull' types is uniformly covered with small, dimple-like indentations and bits of adhering material. These indentations are also visible on 'Shiny' types, but the surface is virtually free of adhering material. At higher SEM magnifications, the surface of 'Dull' types appears rough and ragged whereas the 'Shiny' seeded soybeans have a relatively smooth and undulating surface.

-71-

Without wishing to be bound by theory, it appears that the adhering material on the 'Dull' seeded types are remnants of the membranous endocarp tissue and is rich in hydrophobic protein. The expression of the hydrophobic protein in the endocarp causes bits of this tissue to stick to the seed surface, resulting in the 'Dull' phenotype. Lack of expression similarly may result in the 'Shiny' phenotype. The hydrophobic protein may be involved in the adherence of the endocarp to the seed surface.

Analysis of 'Dull' and 'Shiny' Seeded Varieties

Total genomic DNA was extracted from 'Dull' or 'Shiny' seeded varieties and amplified by PCR using primers targeted to the *HP* gene. The resulting amplification products were clearly polymorphic between the two genotypes. Good amplification of target segments of DNA were regularly observed when template DNA was from 'Dull' types whereas DNA from 'Shiny' types produced multiple products or products that were shorter or longer than expected, depending on the primer combination. These results indicate that different alleles the *HP* gene occurs in 'Dull' and 'Shiny' types of soybean. This allelic variation causes profound differences in seed surface morphology and composition.

To compare *HP* gene structure in two different seed luster phenotypes that were also different in the amount of HP present on the seed surfaces, we hybridized genomic DNA blots with probes derived from the *HP* cDNA sequence under high stringency conditions.

Soybean genomic DNA was isolated from frozen, lyophilized tissue according to the method of Dellaporta et al., (1983). Restriction enzyme digestion of 30 μ g DNA, separation on 0.5 % agarose gels and blotting to nylon membranes followed standard protocols (Sambrook et al., 1989). Digoxigenin labelled cDNA was prepared and used to probe DNA blots according to the instructions provided by the manufacturer (Boehringer Mannheim).

-72-

Hybridization was carried out at 65 °C for 16 h in 0.25 M Na₂HPO₄ (pH 7.2), 20% SDS, 1 mM EDTA and 0.5% blocking reagent (Boehringer Mannheim). Filters were then washed 4 x 15 min at 22 °C in high stringency wash solution (20 mM Na₂HPO₄ (pH 7.2), 1% SDS and 1 mM EDTA), followed by 3 x 15 min washes in the same solution at 68 °C.

A typical result from such a Southern analysis is shown in Figure 18. Genomic DNA blots from cultivars that accumulated large amounts of HP on the seed surface produced strong hybridization signals. These intensely hybridizing fragments are not present in genomic DNA from plants that have only trace amounts of HP on the seed surface. However, several fainter signals are also present in DNA blots from both types of plants. These results indicate that sequences related to the HP cDNA are prevalent in the soybean genome, and that the *HP* gene structure is polymorphic among soybean cultivars. Soybean types that accumulate large amounts of HP on the seed surface possess additional copies of this gene.

All scientific publications and patent documents are incorporated herein by reference.

The present invention has been described with regard to preferred embodiments. However, it will be obvious to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as described in the following claims.

WE CLAIM:

1. An isolated genomic DNA sequence, differentially expressed in seed coat tissues.
2. The genomic DNA sequence of claim 1 differentially expressed within the outer integument of the seed coat.
3. The genomic DNA sequence of claim 1 differentially expressed within the inner integument of the seed coat.
4. The genomic DNA sequence of claim 1 differentially expressed within the thick walled parenchyma of the seed coat.
5. The genomic DNA sequence of claim 1 differentially expressed within the thin walled parenchyma of the seed coat.
6. The genomic DNA sequence of claim 1 differentially expressed within the endothelium of the seed coat.
7. The genomic DNA sequence of claim 1 differentially expressed within the hourglass cells of the seed coat.
8. The genomic DNA sequence of claim 1 differentially expressed within the palisade of the seed coat.
9. The genomic DNA sequence of claim 1 differentially expressed within the stellate parenchyma of the seed coat.
10. The genomic DNA sequence of claim 1 differentially expressed within the membranous endocarp associated with the seed coat.
11. A seed-coat promoter obtained from the genomic DNA sequence of claim 1.

-74-

12. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the outer integument of the seed coat.
13. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the inner integument of the seed coat.
14. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the thick walled parenchyma of the seed coat.
15. The seed-coat promoter of claim 11 that provides for differential expression of a gene associated therewith, within the thin walled parenchyma of the seed coat.
16. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the endothelium of the seed coat.
17. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the hourglass cells of the seed coat.
18. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the palisade of the seed coat.
19. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the stellate parenchyma the seed coat.
20. The seed-coat promoter of claim 11 that controls the differential expression of a gene associated therewith, within the membranous endocarp associated with the seed coat.
21. The isolated genomic DNA of claim 1 characterized by the restriction map selected from the group consisting of Figure 11 (a), (b), (c) and (d).
22. An isolated promoter differentially expressed in seed-coat tissues.
23. The promoter of claim 22 obtained from angiosperms.


-75-

24. The promoter of claim 23 obtained from the group consisting of tobacco or soybean.
25. A cloning vector comprising a heterologous gene encoding a protein, and the promoter of claim 22, wherein the heterologous gene is under the control of the promoter.
26. A plant cell which has been transformed with a vector as claimed in claim 25.
27. A transgenic plant cell containing a promoter as claimed in claim 22, operatively linked to a heterologous gene encoding a protein.
28. A seed containing a promoter as claimed in claim 22, operatively linked to a heterologous gene encoding a protein.
29. An isolated genomic DNA sequence, preferentially expressed in seed coat tissues.
30. A seed-coat promoter obtained from the genomic DNA sequence of claim 29.
31. The seed-coat promoter of claim 11 comprising at least 10 contiguous nucleotides of nucleotides 1-2526 of SEQ ID NO:7.
32. The seed coat promoter of claim 31 comprising nucleotides 1-2526 of SEQ ID NO:7, or an analogue thereof, wherein said analogue hybridizes to a nucleic acid defined by nucleotides 1-2526 of SEQ ID NO:7 under stringent hybridization conditions and maintains seed-coat, or seed-coat associated promoter activity.
33. The seed-coat promoter of claim 11 comprising at least 10 contiguous nucleotides of nucleotides 1-2450 of SEQ ID NO:8.
34. The seed coat promoter of claim 33 comprising nucleotides 1-2450 of SEQ ID NO:8, or an analogue thereof, wherein said analogue hybridizes to a nucleic acid defined by nucleotides 1-2450 of SEQ ID NO:8 under stringent hybridization conditions and maintains seed-coat, or seed-coat associated promoter activity.

-76-

35. The seed-coat promoter of claim 11 comprising at least 10 contiguous nucleotides of nucleotides 1-5514 of SEQ ID NO:9.
36. The seed coat promoter of claim 35 comprising nucleotides 1-5514 of SEQ ID NO:9 or an analogue there, wherein said analogue hybridizes to a nucleic acid defined by nucleotides 1-5514 of SEQ ID NO:9 under stringent hybridization conditions and maintains seed-coat, or seed-coat associated promoter activity.
37. A cloning vector comprising a heterologous gene encoding a protein, and the promoter of any one of claims 32, 34 or 36 wherein the heterologous gene is under the control of the promoter.
38. A plant cell which has been transformed with a vector as claimed in claim 37.
39. A transgenic plant cell containing a promoter as claimed in claim 38, operatively linked to a heterologous gene encoding a protein.
40. A seed containing a promoter as claimed in any one of claims 32, 34 or 36, operatively linked to a heterologous gene encoding a protein.

1/44

**FIGURE 1****SUBSTITUTE SHEET (RULE 26)**

WEST

2/44




FIGURE 2

SUBSTITUTE SHEET (RULE 26)

3/44




FIG. 3B

FIG. 3A

SUBSTITUTE SHEET (RULE 26)

WEST

4/44




FIGURE 4

SUBSTITUTE SHEET (RULE 26)

WEST

5/44




FIG. 5A




FIG. 5B

SUBSTITUTE SHEET (RULE 26)

6/44




FIGURE 6

SUBSTITUTE SHEET (RULE 26)

7144





FIGURE 7

SUBSTITUTE SHEET (RULE 26)

**FIG. 8**

SUBSTITUTE SHEET (RULE 26)

9/44






FIGURE 9

SUBSTITUTE SHEET (RULE 26)

WEST

10/44



SUBSTITUTE SHEET (RULE 26)

WEST

11/44




FIG. 10E

SUBSTITUTE SHEET (RULE 26)

12/44




FIG. 10F





FIG. 10G

SUBSTITUTE SHEET (RULE 26)

13/44

RESTRICTION MAP OF SC20



1 kb

FIGURE 11(A)

SUBSTITUTE SHEET (RULE 26)

WEST

RESTRICTION MAP OF SC21




FIGURE 11(B)

SUBSTITUTE SHEET (RULE 26)

- ☐ BamH I - Restriction Map (1 cut)
- ☐ Bgl II - Restriction Map (1 cut)
- ☐ Hind III - Restriction Map (1 cut)
- ☐ Xba I - Restriction Map (4 cuts)

- ☐ Hps Open Reading Frame:
- ☐ TATA-BOX(1) - Signals
- ☐ Coverage - Direct Repeats





FIGURE 11(C)

SUBSTITUTE SHEET (RULE 26)

16/44



Restriction map of sc4

The shaded and open boxes represent exons and introns respectively.

FIG. 11(D)

17/44

6 DAF




FIGURE 12(a)

SUBSTITUTE SHEET (RULE 26)

WEST

18/44

12 DAF




FIGURE 12(b)

SUBSTITUTE SHEET (RULE 26)

WEST

19/44

18 DAF




FIGURE 12(c)

SUBSTITUTE SHEET (RULE 26)

20/44

FIG. 13A




FIG. 13B




FIG. 13C



FIG. 13D



SUBSTITUTE SHEET (RULE 26)

21/44

FIG. 13E




FIG. 13F





FIG. 13G



SUBSTITUTE SHEET (RULE 26)

WEST

22/44




FIG. 14A




FIG. 14B

SUBSTITUTE SHEET (RULE 26)

23/44




FIG. 14C



FIG. 14D

SUBSTITUTE SHEET (RULE 26)

WEST

24/44

TAAGCTTTCAGAGACAACTGCTTTGAAAAATGGGATCCAAGGTTGTTGCATCCGTTGC 60
M G S X V V J S V J 10

CCTTCTCCTCTCCATCAACATTCTTTTCATTTCATGGTTAGCTCCAGCAGCCACTACGA 120
L L L S I N I L F I S M V S S S S H Y D 30

TCCACAGCCCCAACCTTCTCAGTCACCTGCTCTTATTACAGACCTAGTTGTCCGGATCT 190
P Q P Q P S H V T A L I T R P S C P D L 50

GAGTATTTGCCTCAATATTTTAGGCGGGTCTCTAGGAACCGTGGATGATTGTTGTGCCCT 240
S I C L N I L G G S L G T V D D C C A L 70

CATCGGTGGTCTTGGTGACATTGAAGCCATTGTGTGCCTTTGCATCCAACCTCAGGGCCCT 300
I G G L G D I E A I V C L C I Q L R A L 90

CGGAATATTAAACCTTAACCGTAATTTGCAGTTAATATTAAACCTCTGTGGACGAAGCTA 360
G I L N L N R N L Q L I L N S C G R S Y 110

CCCGTCAAACGCCACTTGCCCCCGAACCTAAGAACAGAAATATGTATGGCACTAATTACCA 420
P S N A T C P R T * 119

TATTACTTCGTATCATGGTGTGTGTTGTTGTCTGTGTTTAAAGTTAAGGATGTTATAC 480

CCTTCGTGCCTGCTACATATATATAGTGGGCACTATAATATTACCAATAAATTAACGTCC 540

ATATATAGAATAATAATAAATAAATAAATATTTCTATACAAATAAAGGTTACGTAATGT 600

TGTTGTTCTCGTGGATGGGGATCTTAICTTCCTCCTCGCTATCTTTGTTATCGTATTTC 660

AGTGAAAGTTGTTCAATAAAAGTCCTTTGTTCAACAAGT (A) 700

FIGURE 15(A)

SUBSTITUTE SHEET (RULE 26)

WEST

25/44

MGSKVVASVALLSINILFISMVSS 25
 SSHYDPQPQPSHVTALITRPSCPD 50
 SICLNILGGSLSGTVDCCALIGGLG 75
 DIEATVCLCIQLRALGILNLRNLQ 100
 LIINSCGRSYPSNATCPRT 119

FIG. 15(B)




FIG. 15(C)




FIG. 15(D)

FIG. 15

SUBSTITUTE SHEET (RULE 26)

WEST

26/44

CULTIVAR: CLARK
PHENOTYPE: DULL

WILLIAMS 82
SHINY



FIG. 16A




FIG. 16B





FIG. 16C

SUBSTITUTE SHEET (RULE 26)

27/44




FIG. 17A




FIG. 17B




FIG. 17C

SUBSTITUTE SHEET (RULE 26)

28/44




FIG. 18

SUBSTITUTE SHEET (RULE 26)

WEST

29/44

```

--C AAT GCT GCG TTA ACT CCT AGA CAT TAC TGG GAA ACG ATG CTT CCA AGA ACT CCC 55
  N A A L T P R H Y W E T M L P R T P 18

TTG CCG AAA GCA ATC ACA GAG CTA CTA AGC CTT GAA AGT AGG TCC ATA TTT GAA TAT 112
  L P K A I T E L L S L E S R S I F E Y 37

GCC GGG AAT GAT GAC CAG TCA GAA AGT AGG TCC ATA TTA GGA TAC GCT GGC TAT AAT 169
  A G N D D Q S E S R S I L G Y A G Y N 56

CAA GAC GAG GAT GAT GTG AGC AAA CAC AAT ATA CAA ATC TTC AAC AGG TTG TTT TTC 226
  Q D E D D V S K H N I Q I F N R L F F 75

TTG GAA GAG GAC CTG CGT GCT GGC AAA ATA TTC AAC ATG AAG TTC GTC AAC AAC ACA 283
  L E E D L R A G K I F N M K F V N N T 94

AAA GCC ACA GTC CCG TTG CTA CCG CGC CAA ATT TCG AAA CAA ATA CCG TTC TCA GAA 340
  K A T V P L L P R Q I S K Q I P F S E 113

GAT AAA AAG AAG CAA GTG TTG GCG ATG CTT GGC GTG GAA GCG AAC TCA AGC AAC GCC 397
  D K K K Q V L A M L G V E A N S S N A 132

AAG ATC ATA GCG GAG ACC ATT GGT CTT TGC CAA GAG CCT GCA ACG GAG GGA GAA AGG 454
  K I I A E T I G L C Q E P A T E G E R 151

AAA CAC TGC GCG ACT TCG TTG GAG TCC ATG GTT GAT TTC GTC GTT TCC GCG CTC GGG 511
  K H C A T S L E S M V D F V V S A L G 170

AAG AAC GTT GGT GCT TTC TCA ACA GAG AAA GAA AGG GAA ACT GAG TCT GGA AAG TTT 568
  K N V G A F S T E K E R E T E S G K F 189

GTA GTG GTG AAA AAT GGG GTG AGG AAG TTG GGA GAT GAT AAG GTT ATT GCC TGT CAT 625
  V V V K N G V R K L G D D K V I A C H 208

CCA ATG AGT TAC CCT TAT GTT GTG TTT GGG TGT CAT CTA GTG CCA AGG AGT AGC GGG 682
  P M S Y P Y V V F G C H L V P R S S G 227

TAT TTG GTG CGC TTG AAG GGA GAA GAT GGG GTT CGA GTG AAA GCA GTT GTT GCG TGC 739
  Y L V R L K G E D G V R V K A V V A C 246

CAC AGA GAC ACG TCA AAG TGG GAC CAT AAT CAT GGG GCA TTC AAA GTG CTC AAT CTT 796
  H R D T S K W D H N H G A F K V L N L 265

AAG CCT GGG AAT GGT ACA GTA TGC CAT GTC TTC ACT GAG GGG AAT CTT CTT TGG CTT 853
  K P G N G T V C H V F T E G N L L W L 284

CCA AAT TAG attaatattaccatatacatattgtccttgttctatccttaataaagtggaatcacctgaagaa 925
  P N * 286

ttgtgcgtaatgagttgtttgtctttgtggaaattgttatctgttgcacaccaaataaggtatatataaaata 1000

acaggagcgtggtattttgttcacaaaaatggatttcaaccgatcaaaaaatatagcctttaccaattagaagg 1075

gtttggcctttgttagcaaaataataaaaaataatatcttgatgg (a)n 1119

```

FIG. 19(A)

SUBSTITUTE SHEET (RULE 26)

WEST

30/44

A

SC4c	FFLEEDLRAG	KIFNMKFVNN	--TKATVPLL	PRQISKQIPF	SEDKKKQVLA	MLGVEANSSN	131
RD22	FFLEKDLVRG	KEMNVRPNAE	DGYGGKTAFL	PRGEAETVPF	GSEKPFSETLK	RFSVEAGSEE	235
PG1B	FFREKMLKSG	TIMPMPDIK-	-DKMPKRSFL	PRVIASKLPF	STSKIAELKK	IFHAGDESQV	472
Sal13-2	FFYKEDLHPG	KTMKVQFTKR	-----	-----PY	AQPY--GVYT	WLTDIKDTSK	215
USP	FF-EHDLHPG	KNFNLGHTNS	VGSIIIR----	-----PF	TKSR--QGVY	--DSIWLANK	111
ADR6	FFYKEDLHPG	KTMKVQFSKP	-----	-----PF	QQPW--GVGT	WLKEIKDTTK	111
SC4c	AKIIAETIGL	CQE-PATEGE	RKHCATSLES	MVDFVVSALG	KN-VGAFSTE	KERETESGK-	188
RD22	AEMMKKTIEE	C-EARKVSGE	EKYCATSLES	MVDFSVSKLG	KYHVRAVSTE	VAKKNAPMQK	294
PG1B	EKMIGDALSE	C-ERAPSAGE	TKRCVNSAED	MIDFATSVLG	RN-VVVRTTE	DTKGSNGNIM	530
Sal13-2	EGYSFEEI--	CIKKEAFEGE	EKFCAKSLGT	VIGFAISKLG	KN-IQVLSSS	FVNKQE----	168
USP	EKQSLEDF--	CYSPTAI-AE	HKHCVSSLKS	MIDQVISHFG	STKIKAISSN	FAPYQD----	164
ADR6	EGYSFEEL--	CIKKEAIEGE	EKFCAKSLGT	VIGFAISKLG	KN-IQVLSSS	FVNKQD----	164
SC4c	FVVVKNGVRK	LGDDKVIACH	PMSYPYVVFG	CHLVPR-SSG	YLVRLKGEDG	VR-VKAVVAC	246
RD22	YKIAAAGVKK	LSDDKSVVCH	KQKYPFAVFG	CHKAMM-TTV	YAVPLEGENG	MR-AKAVAVC	352
PG1B	I-GSVKGING	GKVTKSVSCH	QTLYPYLLYY	CHSVPKVRVY	EADILDPNK	VKINHGVAIC	589
Sal13-2	-QYTVGEVQN	LG-DKAVMCH	GLNFRTAVFY	CHKV-RETTA	FVVPLVAGDG	TK-TQALAVC	224
USP	-QYVVEDVKK	VG-DNAVCMCH	RLNFEKVVFN	CHQV-RDTTA	YVVSILVASDG	TK-TKALTVC	220
ADR6	-QYTVGEVQN	LG-DKAVMCH	RLNFRTAVFY	CHEV-RETTA	FMVPLVAGDG	TK-TQALAVC	220
SC4c	HRDTSKWDHN	HGAFKVLNLK	PGNGTVCHVF	TEGNLLWLPN	*		286
RD22	HKNTSAWNP	HLAFKVLKVK	PGTVPVCHFL	PETHVVWFSY	*		392
PG1B	HVDTSWGP	HGAFVALGSG	PGKIEVCHWI	FENDMTWAIA	D*		630
Sal13-2	HSDTSGMNH-	HILHELMGVD	PGTNPVCHFL	GSKAILWVPN	ISMDTAYQTN	VVV*	276
USP	HHDTRGMNP-	ELLYEALVET	PGTVPVCHFI	GNKAAAWVPN	HTADNLCVM*		268
ADR6	HSNTSGMNH-	QMLHQLMGVD	PGTNPVCHFL	GSKAILWVPN	LSVDTAYQTN	IVA*	272

B


SC4c	--NAALTPRL	YWETMLPRT	LPKAITELLS	L	29
RD22	AIAADLTPER	YWSTALPNT	IPNSLHNL	F	48
Sal13-2	HVHASLPEED	YWEAVWPNT	IPALRDVLK	P	53
USP	GITATSSGED	YWQSIWPNT	LPKTFSDLLI	P	48
ADR6	ARESHARDED	FWHAVWPNT	IPSSLRDLK	P	49

FIG. 19(B)

SUBSTITUTE SHEET (RULE 26)

WEST

31/44



HYDROPHOBIC REGION: [Pattern: Dotted]
CONSERVED SEGMENT: [Pattern: Diagonal lines]
REPEATED REGION: [Pattern: Horizontal lines]
BURP DOMAIN: [Pattern: Stippled]
PEPTIDE CLEAVAGE POINT: ↓

FIG. 19C

SUBSTITUTE SHEET (RULE 26)

WEST

32/44




Fig. 20 C

Fig. 20 B

Fig. 20 A

SUBSTITUTE SHEET (RULE 26)

33/44




FIG. 2/B




FIG. 2/A

SUBSTITUTE SHEET (RULE 26)

WEST

34/44




Fig. 22 A

SUBSTITUTE SHEET (RULE 26)

WEST

35/44




Fig. 22 B

SUBSTITUTE SHEET (RULE 26)

WEST

36/44




FIG. 23A

SUBSTITUTE SHEET (RULE 26)

37/44

caaagttttaac ATG AAA GGC AAT AAT ACA CTT TTG TTG CAT TTA TTC TAC ACT ACT CTC 60
 M K G N N T L L L H L F Y T T L 16

TTC CTG TTT CTT GTA GTG TCA AGT TCA TCT TCA ACA GGG AAT GAA AGT AAC GAT GAC 117
 F L F L V V S S S S S T G N E S N D D 35

ACT AAC AGT AAA GAA GTT TAT ATC GTG TAC ATG GGA GCT GCA GAT TCA ACA AAA GCT 174
 T N S K E V Y I V Y M G A A D S T K A 54

TCT CTT AAA AAT GAG CAC GCT CAG ATT CTG AAT TCA GTG CTA AGA AGG AAT GAG AAT 231
 S L K N E H A Q I L N S V L R R N E N 73

GCC CTA GTA CGG AAC TAC AAG CAT GGT TTC TCT GGG TTC GCA GCT CGT CTA TCA AAA 288
 A L V R N Y K H G F S G F A A R L S K 92

GAG GAG GCA AAC TCA ATT GCT CAG AAA CCT GGT GTG GTG TCT GTT TTC CCT GAC CCC 345
 E E A N S I A Q K P G V V S V F P D P 111

ATT CTG AAG CTC CAC ACT ACA CGT TCA TGG GAT TTC CTC AAA AGC CAA ACT CGT GTC 402
 I L K L H T T R S W D F L K S Q T R V 130

AAT ATC GAC ACC AAA CCA AAT ACG CTG TCC GGT TCT TCT TTT TCT TCA TCA GAC GTC 459
 N I D T K P N T L S G S S F S S S D V 149

ATT CTT GGC GTC TTA GAC ACA GGC ATA TGG CCA GAG GCG GCG AGT TTT AGC GAC AAG 516
 I L G V L D T G I W P E A A S F S D K 168

GGT TTC GGT CCT GTT CCA TCC CGA TGG AAA GGC ACC TGC ATG ACA TCA AAA GAC TTC 573
 G F G P V P S R W K G T C M T S K D F 187

AAT TCC TCT TGT TGT AAC AGG AAG ATA ATT GGC GCG AGG TTT TAC CCT AAC CCA GAG 630
 N S S C C N R K I I G A R F Y P N P E 206

GAG AAA ACG GCA AGG GAT TTC AAC GGA CAT GGG ACT CAC GTT TCG TCG ACG GCA GTG 687
 E K T A R D F N G H G T H V S S T A V 225

GGC GTG CCG GTG AGT GGC GCA TCG TTC TAT GGT CTG GCG GCG GGG ACG GCA AGG GGT 744
 G V P V S G A S F Y G L A A G T A R G 244

GGG TCC CCT GAG TCA AGG TTG GCG GTT TAC AAA GTG TGT GGG GCT TTT GGG TCA TGT 801
 G S P E S R L A V Y K V C G A F G S C 263

CCT GGG TCG GCC ATT CTT GCG GGG TTT GAC GAT GCC ATT CAC GAC GGA GTG GAT ATC 858
 P G S A I L A G F D D A I H D G V D I 282

TTG TCG CTG TCG CTC GGT GGA TTC GGT GGA ACC AAA ACC GAT TTG ACC ACC GAC CCG 915
 L S L S L G G F G G T K T D L T T D P 301

ATT GCG ATT GGA GCA TTC CAC TCC GTC CAG CGC GGC ATC CTG GTG GTC TGC GCC GCC 972
 I A I G A F H S V Q R G I L V V C A A 320

GGG AAC GAC GGA GAA CCA TTC ACC GTT CTC AAC GAC GCA CCT TGG ATT TTA ACC GTT 1029
 G N D G E P F T V L N D A P W I L T V 339

GCA GCT TCC ACC ATC GAC CGT GAT CTT CAA TCC GAC GTG GTC TTG GGT AAT AAC CAA 1086
 A A S T I D R D L Q S D V V L G N N Q 358

FIG. 23(B)

SUBSTITUTE SHEET (RULE 26)

38/44

GTC GTC AAG GGA AGA GCC ATA AAT TTC TCC CCT CTT TTA AAT TCT CCC GAT TAT CCA 1143
 V V K G R A I N F S P L L N S P D Y P 377
 ATG ATA TAT GCT GAG TCT GCT GCC AGG GCA AAT ATC TCC AAC ATA ACT GAT GCA AGA 1200
 M I Y A E S A A R A N I S N I T D A R 396
 CAA TGC CAC CCA GAT TCA TTA GAT CCA AAA AAA GTC ATA GGG AAG ATT GTG GTT TGT 1257
 Q C H P D S L D P K K V I G K I V V C 415
 GAT GGA AAA AAT GAC ATT TAT TAT TCA ACT GAT GAG AAA ATT GTC ATA GTG AAG GCG 1314
 D G K N D I Y Y S T D E K I V I V K A 434
 TTG GGA GGA ATA GGT CTG GTT CAT ATT ACT GAT CAA TCT GGA TCA GTA GCA TTT TAT 1371
 L G G I G L V H I T D Q S G S V A F Y 453
 TAT GTG GAC TTC CCA GTA ACA GAG GTA AAA TCA AAA CAT GGC GAC GCA ATC CTC CAG 1428
 Y V D F P V T E V K S K H G D A I L Q 472
 TAC ATC AAC TCA ACT AGC CAT CCA GTG GGA ACA ATA CTA GCA ACA GTT ACA ATT CCT 1485
 Y I N S T S H P V G T I L A T V T I P 491
 GAT TAT AAG CCT GCT CCC CGG GTG GGT TAT TTT TCA TCA AGA GGG CCT TCA TTG ATT 1542
 D Y K P A P R V G Y F S S R G P S L I 510
 ACA AGC AAT GTT CTC AAG CCT GAT ATT GCA GCC CCG GGA GTT AAC ATT CTC GCT GCA 1599
 T S N V L K P D I A A P G V N I L A A 529
 TGG TTT GGA AAT GAC ACA TCA GAG GTT CCA AAA GGA AGA AAG CCC TCA CTA TAT CGC 1656
 W F G N D T S E V P K G R K P S L Y R 548
 ATA CTC TCA GGA ACT TCC ATG GCT ACT CCA CAT GTT TCA GGG CTT GCA TGC AGT GTC 1713
 I L S G T S M A T P H V S G L A C S V 567
 AAA AGA AAA AAC CCC ACT TGG AGT GCC TCC GCA ATC AAA TCT GCC ATC ATG ACT TCA 1770
 K R K N P T W S A S A I K S A I M T S 586
 GCA ATT CAA AAT GAC AAT TTG AAG GGT CCC ATA ACA ACG GAT TCA GGG TTG ATA GCC 1827
 A I Q N D N L K G P I T T D S G L I A 605
 ACA CCT TAT GAC TAT GGA GCA GGG GCA ATT ACA ACA TCT GAA CCA TTG CAA CCG GGG 1884
 T P Y D Y G A G A I T T S E P L Q P G 624
 CTA GTT TAT GAA ACC AAC AAC GTT GAC TAC TTG AAC TAT TTG TGT TAC AAT GGA CTT 1941
 L V Y E T N N V D Y L N Y L C Y N G L 643
 AAC ATA ACC ATG ATA AAG GTC ATC TCC GGA ACT GTC CCC GAG AAT TTC AAT TGT CCC 1998
N I T M I K V I S G T V P E N F N C P 662
 AAG GAT TCG AGC TCT GAT CTC ATC TCC AGC ATC AAC TAC CCT TCC ATA GCA GTA AAC 2055
 K D S S S D L I S S I N Y P S I A V N 681
 TTC ACT GGC AAA GCA GAC GCG GTC GTG AGT AGA ACT GTC ACA AAC GTT GAC GAA GAA 2112
E T G K A D A V V S R T V T N V D E E 700
 GAT GAA ACA GTG TAC TTC CCC GTT GTT GAA GCT CCT AGT GAA GTA ATT GTC ACA CTC 2169
 D E T V Y F P V V E A P S E V I V T L 719
 TTT CCA TAT AAT CTT GAG TTT ACG ACA AGT ATT AAA AAA CAA AGC TAC AAT ATT ACT 2226
 F P Y N L E P T T S I K K Q S Y N I T 738

FIG. 23(B)(Cont'd)

SUBSTITUTE SHEET (RULE 26)

WEST

39/44

TTC AGA CCG AAG ACC TCC TTG AAG AAA GAT TTG TTT GGA TCT ATC ACT TGG AGT AAC 2283
E R P K T S L K K D L F G S I T W S N 757

GAC AAA TAT ATG GTT CGA ATT CCT TTT GTA TTA ACT AAA TAG tgaaattaaaaagtagcga 2344
D K Y M V R I P F V L T K * 770

tgaaataaatgcaagctaagttcttctgtggtgcctacactcgagtcctgattatttattattccatagccttctgt 2419

tttaatttat ttattatact ttcagcct(a)n 2447

FIG. 23(B)(Cont'd)

SUBSTITUTE SHEET (RULE 26)

WEST

40/44




FIG. 23C

SUBSTITUTE SHEET (RULE 26)

WEST

41/44

D region

SC20:2
AF70
Cucumisin
P69B
Ag12
Subtilisin BPNE
Kex2
Furin

*
SDVILGVLDGTGI 156
TDIILGFLDTGI 145
SNIVVGVLDTGI 143
KGVIIIGVIDTGI 149
EDVIIIGVIDSGV 148
SNVKVAVIDSGI 142
AGVVAAIVDDGL 178
HGIVVSILDDGI 156

H region

SC20:2
AF70
Cucumisin
P69B
Ag12
Subtilisin BPNE
Kex2
Furin

*
DFNGHGTHVSSTAVG 224
DYQGHGTYTAATAAG 229
DNTGHGTHTASTAAG 214
DDIGHGTHTASTAAG 213
DTLGHGTHTASTAAG 216
DNNSHGTHVAGTVAA 181
SDDYHGTRCAGEIAA 223
NDNRHGTRCAGEVAA 204

S region

SC20:2
AF70
Cucumisin
P69B
Ag12
Subtilisin BPNE
Kex2
Furin

*
SGTSMATPHVSGLA 562
SGTSVAVPHVTGAA 571
SGTSMSCPHITGIA 535
SGTSMSCPHLSGVA 541
SGTSMACPHASGVA 547
NGTSMASPHVAGAA 338
GGTSAAAPLAAGVY 395
TGTSASAPLAAGII 378

N region

SC20:2
AF70
Cucumisin
P69B
Ag12
Subtilisin BPNE
Kex2
Furin

SVQRGILVVCAAGNDG 322
ATQKGILVVSSAGNEG 329
AVERGILTSNSAGNGG 310
ATERGILVSCSAGNSG 308
AMEKGVVSTTSAGNAG 318
AVASGVVVVAAAGNEG 264
RDSKGAIYVFASGNGG 316
RGGLGSIPVWASGNGG 297

FIG. 23(D)

SUBSTITUTE SHEET (RULE 26)

WEST

42/44




FIG. 24A



FIG. 24B

SUBSTITUTE SHEET (RULE 26)

WEST

43/44




Fig. 25 A

Fig. 25 B

SUBSTITUTE SHEET (RULE 26)

WEST

44/44




Fig. 25 C

SUBSTITUTE SHEET (RULE 26)

WEST

1/33

SEQUENCE LISTING

<110> Miki Dr., Brian L
Gizen Dr., Matk
Miller Dr., Shea
Boutilier Dr., Kim
Hu., Ming
Bowman, LuAnne
Batchelor, Anthea

<120> Seed-Coat Promoters, Genes and Gene Products

<130> 08-869947WO

<140>

<141>

<150> 09/059,090

<151> 1998-04-13

<160> 9

<170> PatentIn Ver. 2.0

<210> 1

<211> 1070

<212> DNA

<213> Nicotiana tabacum

<400> 1

tctagacttg tcttttcttt acataatcct cttcttcttt tttttgtag tttctctgt 60

tttatccaaa aaacgaatta ttgattaaga aatacaccag acaagttttt tacttctttt 120

tctttttttt tttgtggttaa aaaattacac ctggacaagt ttatcacgaa aatgaaaatt 180

gctattttaag ggatgtagtt ccggactatt tggaagataa gtgtaacaa aataaataaa 240

SUBSTITUTE SHEET (RULE 26)

WEST

2/33

taaaaagttt atacagttag atctctctat aacagtcac cttatttata acaatacttt 300
actataaccg tcaaatttat ttgaaacaa aattttcatg ttatgttact ataacagtat 360
tttattatag caaccacaaa atatcgaaac agatacgatt gttatagagc gatttgattg 420
tattcattatc cacatatattt cgtaagccca attactctc ctacgtacga tgaaagtaaa 480
ccaatttaaa gttgcaaaaa tocaatagat ttcaatactt cttcaactgg cgttatgtta 540
ggtaatgact cttttttaac ttttcatctt taatttgaag tttctttcat taaaagaaa 600
tttctagaag agaagtgtt taacacttct agctctacta ttatctgtgt ttctagaaga 660
aaaatagaaa atgtgtccac ctcaaaaaca actaaagggt ggcaaactc cacctattta 720
ttttattttg gattaattaa gatatagtaa agatcagtta taaacggagt ttgagttga 780
tacagtgaat tttaagatgt gtaccgattt aactttattt acatttatgt ttgcacata 840
taagaagtcc gatttggaaa tactagattt tgtcaatcag gcaattcatg tggttgaaga 900
atttaagtta tatacaatga tgatataaag aatttttata ctattagtgc aaattaatcg 960
attactaaaa attattattc tattaattta tgctatcgtg cctccccaac cgtcgaccg 1020
cggtaaccgg tggtcagtc cttatgttac gtctgtaga aaccccaacc 1070

<210> 2

<211> 4700

<212> DNA

<213> Glycine max

SUBSTITUTE SHEET (RULE 26)

WEST

3/33

<400> 2

tagataaaaa aatgggatat aatTTTTtctc agatgttggt tatactgttt ttttaaatcag 60

aattaaaatt cctctttaat tatcgacata attttttttg gtgaatatta tcgacataat 120

tatttaatac aaatttttat tgtacataga agtgatactt caattttaat attggagaac 180

agtagcaaaa cataaaaaaa ctgttattag aagaaaaaaa tatatggaaa aggttagcta 240

catatattag cttaaattagt tgttctaatt ggctatataa accctattgt actctttgta 300

atctcacctt tttcatttaa atacatttct actttttaag ttctatattt tctctcaatt 360

ttcttcgata aaccatgaaa tttaacatgg tatatcagcg ataccacca ctttgaaagc 420

catgtatggc tagtatgggc agccaaaatt tgccttggtt caagcaaagc aagtgtttat 480

atagatgtga cttttgttga ggaactcatg ccaatggtac tgattgtgaa actgagaaaa 540

ctaatttga gaatttgaat tatgatcatt aaatactcct ctctgacta ccttcgtccc 600

tcaaatttgt accatcatta tttcccaaaa atttgattac aatgcactaa ttaatgaatg 660

tttcttacat tatcatatta tcatatctga cattttgttt ttacttttta taataattat 720

tttaaaaagt catacatgca aataattttt taatagttaa cagttaaatt tttacagtaa 780

aatgcatga aaattaaact ttatttttcc aagtcacat ttagtcaa atccaaaacaa 840

tgattatttt ttgcaaatga atgtttattg aacatttaa tgtagcctaa ttaattctgg 900

ttatggtgtc aatgttccaa aacctaatgc aagatcttag caagtacata catagatcta 960

attttaaact tatctttacg caagagatat aaagattata catctagttt taaacattaa 1020

SUBSTITUTE SHEET (RULE 26)

WEST

4/33

cttttgTTTT tgtgttaaaa aacagtaaca ttttcttaat tttgtagagt gacgtgctcc 1080
aaccatatta acgaagattt taattggtat tcaagttcat gaacttagta aataagtttt 1140
ggtcttcagt tttcaatttt cattacaaca tttatgtaaa atatcaacgt tttctgaaat 1200
ttgttgcttg tgtgctccaa ccacatttaa gagattatag aaattaattt tcaagaagat 1260
aatgattcct actcttgctg gccctaccat agtacaataa atccactcat aaatcaacaa 1320
gtcgtcgtca taggcaattg ggcacatata cataaacaat acgtacgtga tattatctag 1380
tgtctctcag tttactttat gagaaattat ttttctttaa aaaaagttaa ttaataaaaa 1440
catttgcgat accgtgagtt acaagaaatc cgcgaattc atctctataa ataaaaggat 1500
ctatatgaga ggtaaaatca tattaactca aaatgggttc catgctgcta ttagtagtgg 1560
cattgttggt tgcatttget atgcatgcag gtttttcagt ctcttatgct cagcttactc 1620
ctacgttcta cagagaaaca tgtccaaatc tgttccctat tgtgtttgga gtaatcttcg 1680
atgcttcttt caccgatccc cgaatcgggg ccagtctcat gaggttcat tttcatgatt 1740
gctttgttca agtacgtact ttttttttcc cttccaaaat gccctgcata ttaacaaga 1800
ttgctttggt cacctagaaa aatgtgtttt tttcaacgat cttacgtacg tttgtttggt 1860
ttgaaaaata aatcagaaag agatcaagaa aatagctaga aagaaagcaa cgttttttta 1920
aaaggatttt agtgtgagaa aaatattaaa actgaagaga aagaaattaa ataagctttt 1980
cttgaatgat atttacatgt cttattaact taaagtcacc ttttttcttt aagttgtgct 2040

SUBSTITUTE SHEET (RULE 26)

WEST

5/33

tgaagaaaa agatgtcttt cagtttagtt ttgattaatg ctaattatat ttttaattaa 2100
ttaattaata ctatatatct atttaccata ttaattatta ctatatttca tgatgacaac 2160
agacaagtat tctaaagagg tatcggtaga tgattaattt ttttataaaa aaatcttttg 2220
cgtgtataga tattctttta taattggtgc agaaacttgt aatgctaatt gcaattaate 2280
ttacattgat taactaatag ctataatcaa tatttaggtt aggtatagga gacaaatcaa 2340
gtgatctgaa caaattaagt tgttatattt gcattgtgac agggttgtga tggatcagtt 2400
ttgctgaaca aactgatac aatagaaagc gagcaagatg cacttccaaa tatcaactca 2460
ataagaggat tggacgttgt caatgacatc aagacagcgg tggaaaatag ttgtccagac 2520
acagtttctt gtgctgatat tcttctattt gcagctgaaa tagcttctgt tctggtaatt 2580
aataactcct aattaattcc caaccattaa aaagttgcat gattggattc aaaattctat 2640
ggtattgggg ttctgatata aatttgtaat taaattgcac taaaaaaaaat tatcatatac 2700
ttttaataaa aaaaatttat ctaatttaat ttattattaa aactattttt aaaattcaat 2760
cctaactctt ttttaatcgg agcatgtaag ctggcaccca ccgtatatcg ttggaagatg 2820
ctataaaacc atttaattaa tggatggaat cagtcaaaac atttaattca aaatactctt 2880
aattgtgatt agtaatcatg ttggggcaag ttacgttgtg tataattaat ttgacttaat 2940
cagataaaaa aacaaatgga cgcaagccgg ttggtataga tatcactggc ctgtagaata 3000
tgtggttttt cacgttttaa taaaagctag ctactatatt atatttagtc tttttttttc 3060

SUBSTITUTE SHEET (RULE 26)

WEST

6/33

ttaaaccat ttaacgtgat ttattgactg tgaacatgt ttccacacac aggcttagaa 3120
actcctcgca actaacatct ccaaaatttg actattttatt tatgaagata attcatctat 3180
gatgttcaac tctattatat atatgtatca tcgcagtatt aagaattata atagtcaaatt 3240
atagaagtat atcggtgtaa tgtagttgca tgtgcgacct gtttcgtgta aaatgcttat 3300
tctatatagc tttttttatt ggaaaataac gatgaactaa aaacgaaagg gtatcatata 3360
gtttgacttt tatgttagag agagacatct taatttgggc atatgttaa taattaatta 3420
caatgcatac acaaatattt atgccatct taaaaaatga taaaatatca taggtatact 3480
caactatatg atatcccat aacagaaatt gtacttttct tcaggcaatg aacttaacat 3540
ttctgtttgc taaaaacaaa catccactta aagtgggtca acatatttat gtaataattt 3600
acagggagga ggtccaggat ggccagttcc attaggaaga agggacagct taacagcaaa 3660
ccgaaccctt gcaaatcaaa accttccagc acctttcttc aacctcactc aacttaaagc 3720
ttcctttgct gttcaaggtc tcaacacct tgatttagtt acactctcag gtatacataa 3780
tcaatttttt atttgctatt agctagcaat aaaaagtctc tgatacagac atatttagat 3840
aaattaattt ctccataaac atttataata aaattatcaa tttatgtact taaaaattat 3900
ggattgaagc tcttttcac caacttttac taaagttaag gtgcatataa tataaaataa 3960
actatctctt gtttcttata aaaagattga agataagtta aagtctactt ataaatcatt 4020
aatatatgta taggtgggtca tacgtttgga agagctcggt gcagtacatt cataaaccga 4080

SUBSTITUTE SHEET (RULE 26)

WEST

7/33

ttatacaact tcagcaacac tggaaaccct gatccaactc tgaacacaac atacttagaa 4140
gtattgctg caagatgcc ccagaatgca actggggata acctcaccaa tttggacctg 4200
agcacacctg atcaatttga caacagatac tactccaatc ttctgcagct caatggctta 4260
cttcagagtg accaagaact tttctccact cctgggtgctg ataccattcc cattgtcaat 4320
agcttcagca gtaaccagaa tactttcttt tccaacttta gagtttcaat gataaaaatg 4380
ggtaatattg gagtgtgac tggggatgaa ggagaaattc gcttgcaatg taattttgtg 4440
aatggagact cgtttggtt agctagtgtg gcgtccaaag atgctaaaca aaagcttgtt 4500
gctcaatcta aataaaccaa taattaatgg ggatgtgcat gctagctagc atgtaaaggc 4560
aaattagggt gtaaacctct ttgctagcta tattgaaata aaccaaagga gtagtgtgca 4620
tgtcaattcg attttgccat gtacctcttg gaatattatg taataattat ttgaatctct 4680
ttaaggctact taattaatca 4700

<210> 3

<211> 1121

<212> DNA

<213> Glycine max

<400> 3

caatgctgctg ttaactecta gacattactg ggaacgatg cttccaagaa ctccttgcc 60
gaaagcaatc acagagctac taagccttga aagtaggtcc atatttgaat atgccgggaa 120

SUBSTITUTE SHEET (RULE 26)

WEST

8/33

tgatgaccag tcagaaagta ggtccatatt aggatacgtt ggctataatc aagacgagga 180
tgatgtgagc aaacacaata tacaaatctt caacaggttg tttttcttgg aagaggacct 240
gcgtgctggc aaaatattca acatgaagtt cgtcaacaac acaaaagcca cagtcccggt 300
gctaccgcgc caaatttcga aacaaatacc gttctcagaa gataaaaaga agcaagtgtt 360
ggcgtatgctt ggctgtgaag cgaactcaag caacgccaag atcatagcgg agaccattgg 420
tctttgccaa gagcctgcaa cggaggggaga aaggaaacac tgcgcgactt cgttggagtc 480
catggttgat ttcgtcgttt ccgcgtcgg gaagaacgtt ggtgctttct caacagagaa 540
agaaagggaa actgagtctg gaaagtttgt agtggtgaaa aatgggggtga ggaagttggg 600
agatgataag gttattgcct gtcaccaat gagttacct tatgttgtgt ttgggtgtca 660
tctagtcca aggagtagcg ggtatttggt gcgcttgaag ggagaagatg gggttcgagt 720
gaaagcagtt gttgcgtgc acagagacac gtcaaagtgg gaccataatc atggggcatt 780
caaagtgtc aatcttaagc ctgggaatgg tacagtatgc catgtcttca ctgaggggaa 840
tcttctttgg cttccaaatt agattaatta ccatatacat atttgcctt gttctatcct 900
taaataagtg gaatcacctg aagaattgtg cgtaatgagt tgtttgtctt tgtggaaatt 960
gttatctgtc ttgcatcacc aaataggtat atataaata acaggagcgt ggtatttgtt 1020
gcacaaaaat ggatttcaac cgatcaaaaa aatatagcct ttaccaatta gaagggtttg 1080
gctttgttag caaataataa aaataaata tcttgatgg 1119

SUBSTITUTE SHEET (RULE 26)

WEST

9/33

<210> 4

<211> 2723

<212> DNA

<213> Glycine max

<400> 4

taatacgact cactataggg cgaattgggt accgggcccc ccctcgaggt cgactcgatc 60
tcaaatttta ttccatttaa aataaaacat aatttaattt tegtctctct tccttattgt 120
atcattataa aagtaggaaa acaaatataa attagaacaa acataatatt aattaataag 180
aataatttgt ttgttgcttt gaattttcta ttctaataac attaggtagt aataaaatta 240
agttgagttt catttttttg aaagaattaa cttaataatt gtatattttt gtttaagtat 300
aatatttttag ataatgtatt atcacattaa aaatttagag tgatagacaa attatgttta 360
ttaatcaata ttatgtttat ttaattgttt gttttaagtt aggtttgttt tcatattttt 420
tttagtgttt ttattataat gagaaaaaaaa atggaggata aaagataaaa attatattat 480
attttactct taaataaaac aaaatttgga gtccataaaa ttagtttaat gaaaacttgt 540
tgacataggt ctaatctatt caataatcat gttagcttat ttgtgctctt ggactccttc 600
attaacggta atagatgaat gaatttattg catttccttc acttttgatt actaaaattt 660
aaattttcat ttttcgaaaa tgaattcgtt actattttat acatttaaaa ataaaaatga 720
ttatttcac ttataattat caaataatat gcctgtgatg atgatttttt tcaaaaattg 780
aaaatgtctt attgcctagt taggctatag aatctttttg gtcacatca atcgcatggc 840

SUBSTITUTE SHEET (RULE 26)

WEST

10/33

gacatgggta cgcggttcat tgggtaatta tatttatgta tttaaattaa ttctattaaa 900

accaattcta ttaaagctga aaactaaat gacacttaat ttctgtaaga gtcgtgtaat 960

tagcgagcca gagtaatgca accaaagagc ttctttcccc accttaattc ctttataatg 1020

aatttgattg ataccataca tagctaagct tttttttctt tgcatttaca tagccaagct 1080

ttagctgcta tatataatat gttgtttata taattttgac atgttgcatt acatgttact 1140

aactggttgc aacctccagt tttggttacc gaatatatgc tggccggcac caactacaaa 1200

aaatttctgt atggctggct actaatacta tatcttcaaa tgtctcactt tacatggact 1260

gagtgcataa cgggcaattg cagtgtcat aaggaatacg catgcaatca ttttatcatc 1320

gcagattgtt gcttttccct cactgtaaaa aaaaataaaa ataacaatta attgagcacc 1380

gaatgaacta gaccgattgc atgcattata atacatgaag aaaattattt atactaatct 1440

tttgtttcaa gttttaagct tgactttttt gttttacaag cacaataaca atcaaataac 1500

caataattta atcattaaaa agataaaca actcattaga taaaataata attatgataa 1560

aatatgttta agtcctctat atttaagttg aattttgttt ttagtctttc aaattgtgca 1620

ttttagtctc tgaaatttta ttttaaaaaa atagtctttg ggcttactct ccaaatttg 1680

cacattttta ttttaggtta ttgattaaag attaaaaaca ttttttttta tcaaatttag 1740

ggaccaaatt acttgatata aatactcaag aactaaaaac aaatttcaat aaaaatataa 1800

gggattaaaa atatatttta ttctgataac taacaatcat caaattttat gacaatcata 1860

SUBSTITUTE SHEET (RULE 26)

WEST

11/33

tgtaaccttt tataatataa ttgtaagaat aattaatttt aataaataaa taaataaatg 1920
agacttataa tagtaaacat atatgtttgc ctaggtaata acaatgataa caagtaatag 1980
aattatcatt cttattgtat ctattaatta atatattatt agatgtatta attagtatat 2040
atatttttta ctataaatag tatatattat taattattat acatttatta aagtatatcc 2100
attaaagatt ttgagggagg ggagctgaag ccactaaccc ccgtaaatcc gtccttgac 2160
aaaacacgac atgagaatgg ttttgatataa tccacaattt aatatccaat aaaataattt 2220
cttttttatt ttaattagga aactccaaga ttgctttaac ttatacaaaa tctgaataac 2280
acaaaaaat aaaaaatcgg aattaaatgt tgcccgattg ttgactaact tatacaaat 2340
cttattttaa tgettaaat cgtgtcataa tataatgaat atatatttgc aaatatatat 2400
ttattatata attgcaaata tatattctaa ttttgagtat aaataacagc atgtgagggt 2460
gcagcaaac acacactgag tgcaacaaag ttttaacatg aaaggcaata atacactttt 2520
gttgcattha ttctacacta ctctcttctt gtttcttgta gtgtcaagtt catcttcaac 2580
aggaatgaa agtaacgatg acactaacag taaagaagtt tatatcgtgt acatgggagc 2640
tgcagcccg gggtaccact agttctagag cggccgccac cgcggtggag ctccagcttt 2700
tgttcccttt agtgagggtt aat 2723

<210> 5

<211> 829

<212> DNA

SUBSTITUTE SHEET (RULE 26)

WEST

12/33

<213> Glycine max

<400> 5

gccttaaggc aacgacagcg agttcttctg ttgttcgttg actccaagga cggggtctta 60
gttggtggct tcgtggttcc cttcttttggg ggcttcgttg ttgttgtctt tgtcatcctt 120
gttggtgatg tcttcttcgg ttccgtctcc tcggatttct tcaacgtcaa ctctggctcc 180
tccaccactg tttctctgt ctcttctct tcggtcgtcg tcttctcttt agtctctct 240
tttgacttct tcactttctt ctctgtcttc tcggatttct ttgatgactt tgtctttgtc 300
ttagtcgtgg tcgtcgttgg tgtctcttc tcttgtttgg tcgactctgt caccttcggg 360
gtcatctcca aggacaactc ttttgacttc gaatctgaat acctgtcaca ctcaactctt 420
ttgttcaaat ttttaccagg atcaacaccc ctagtaatcc agatggtaaa acgtacagta 480
caacttttctg aaaaaaaaa taaataaaga aatcaaatga aataataatt atataataat 540
aatatactac caattcagaa accaacataa tactctccat atggatgcac actcgtgttg 600
tgaaccgagg ttagtcgca cgaatggtgc accactgtac tctttaagat cctcaacaca 660
taacttcact cgttcgacga cgaccacgac ctcaaatccc tactcccaa taaaataacc 720
gtcccaatgt actcaaacac tactacacaa taacacaaag aaatccaaac ttatttacta 780
taaataaaat tttttttttt ttttttagat ctagctggcg tcgggttga 829

<210> 6

SUBSTITUTE SHEET (RULE 26)

WEST

13/33

<211> 700

<212> DNA

<213> Glycine max

<400> 6

taagctttca agagacaaac tgctttgaaa aatgggategc aaggttggtg catccgttgc 60
ccttctcttc tccatcaaca ttcttttcat ttccatggtt agctccagca gccactacga 120
tccacagccc caaccttctc acgtcactgc tcttattaca cgacctagtt gtccggatct 180
gagtatttgc ctcaatattt taggcgggtc tctaggaacc gtggatgatt gttgtgcctt 240
catcgggtggt cttggtgaca ttgaagccat tgtgtgcctt tgcaccaac tcagggcctt 300
cggaatatta aaccttaacc gtaatttgca gttaatatta aactcctgtg gacgaagcta 360
cccgtaaacc gccacttgcc ccggaacctc agaacagaat atgtatggca ctaattacca 420
tattacttgc tatcatggtg tttgtttggt tgtctgtgtt taaagttaag gatgttatac 480
ccttcgtgcc tgctacatat atatagtggg cactataata ttaccaataa attaacgtcc 540
atatataaga ataataataa ataaataaat atttctatac aaataaaggc tacgtaatgt 600
tggtgttctc gtggatgggg atcttatctt cctcctcgct atctttgttt atcgtatttc 660
agtgaaagtt gttcaataaa agtcctttgt tcaacaagta 700

<210> 7

<211> 3368

SUBSTITUTE SHEET (RULE 26)

WEST

14/33

<212> DNA

<213> Glycine max

<400> 7

tctttcgatc aatactaata aagtcttatt tgccttcag agacaattga gtccgttggc 60
acgcagagac aaattatggt aatttgcccc tttttgaaga cttcaatgct tttcgatcaa 120
gactattaata gtcttctttg ccttctagag acaaattatg gtcctctgat tctttttgaa 180
tacttcaatg tctttcaatc aagacaatca aagttttttc gaatacttca aagtcttctt 240
tgccttcagg agacaattaa gtctgttggg acgcagagac aaattatgat catctacccc 300
atttcgaaga cttcaatgct tttcgataaa gactattaata gtcttctttg ccttcaggag 360
acaatcaagt tcattggcac gtagagacaa attatggtca tctgcctctt ttogaatact 420
tcaatgtctt tcgattaaga ctatcaaagt cttcttttcc ttccggagac aatcagggtc 480
tttggcacgc agagacaaat tatgttcac tgcctctttt cgaagacttc aatgtctttc 540
gatcaagact atcaaagtct tctttgcctt ccgaagacaa tcaagtctgt tggcacgcaa 600
agtttgagga aaaattggac gaagatcggg acaaattggac cgtatggttt gacggagcgt 660
caaacattct aggccatggc attggggcag tattggtctc tccggacaat caatgtgtac 720
ctttcacaac caggctagga ttcgactgca ccaacaacga tcttcggtgc aattactcaa 780
atcggggaac taagggtgaa tgaagttgct gtcatttttg gcaaggacaa gggagttgtt 840
ggagctaacc atggcatagc aaaggagatt gagggagagg ataattaggg catcaacacc 900

SUBSTITUTE SHEET (RULE 26)

WEST

15/33

catggaaccc atttaatttc ccaacataga ttgatagaaa tattattgca gtctctcttg 960
atagcttaaa tattgatgag caagtgtctt tgettgtggt ttctagctga actttacagg 1020
tacgagtata agattactaa acttggttctg atcctgaacc cgaactcgcc tgtcacttaa 1080
aatttttaaa atttttgcat aatttaatca aaaggcataa aatttttatt actagttaat 1140
ttttttttta gaatttttac ataatttaatt attattttct taactatttt ttagatacac 1200
gcgctataat aaatttattt atatatatgt agttaaaaaat aaatgtttta taatcaattt 1260
attttttcta aaatcaaatt tttaattttt ttttataaaa aaatattttt ctaagttgaa 1320
tcgtgtatgg gacggggtca gggatacccg atatctgacg ggtacgagga tgagacaata 1380
aacttaaata cgtcgagtat tggatacgag tatgggaata tgttggggag tcggggtaag 1440
gaattgagga aacaatatcc ataccacccc gccctattgt catgtctaga cactacaaag 1500
aagggttaaa gaaacctaag ttaaaatagt agattatatg acatttagtc ctgtaaaaag 1560
aagaagagaa agatgtagaa aattttcaag aaagatatca agttaataa tatttttcaa 1620
agtttgattt ttaattatat caaacaacgt agtgtgattc atgtaattgg tgacttacct 1680
actagtataa aaatttggtc tctttgttgt tgttgcatgt atggaatgaa ttttaaaaaa 1740
atcataaata taatttgaaa tcattttaaa attatgtaa atcatttcga attattgatc 1800
tagattaaac aattacttag tgtaacaaga gaatttttgc ttagatttaa actttaattc 1860
ggctagcacc tagagattta tttttgtaat gatccatgac aatatcataa ttatgataat 1920

SUBSTITUTE SHEET (RULE 26)

WEST

16/33

atattgcata atttaaattt gtattcatct ttctttaaaa aatatacttg aaagtgttaa 1980
attgtacttc aaagatttag catattagtt tagttctgga taataaatta aaattattat 2040
tctcaaaaat gagataattc tttcatgtac aattcttcat acatagtatc aaatgtcttc 2100
ttcattttat gacacatgcc ttttaatttt tatattaata aaattaattt tttattaaat 2160
taaataatat tttaatctct ttaatgcttg aattaatata ttttttttta aaaaactaag 2220
catgacaagg tatttacaat ttactctaga aataatatac actaattaac acaagaataa 2280
gtatttttca aaatattttt ttttccatc aaaccacaag tatctgcaac aaaacttctt 2340
ttgagtgttt aagagagtta cataccctaa acagaaatgt gggaccgttg atcatcacac 2400
caattcaatt tattcagacg ctgctttgtt ggtaattggc ctataaattg tatcccaaac 2460
ttcagttaga caacaaaagc acttggtcac caattaagct ttcaagagac aaactgcttt 2520
gaaaaatggg atccaagggt gttgcacccg ttgcccttct cctctccatc aacattcttt 2580
tcatttccat ggtagctcc agcagccact acgatccaca gcccacact tctcacgtca 2640
ctgctcttat tacacgacct agttgtccgg atctgagtat ttgcctcaat attttaggcg 2700
ggctcttagg aaccgtggat gattgttgtg cctcatcggg tggctcttgt gacattgaag 2760
ccatttgtg cctttgcac caactcaggg cctcggat attaaacctt aaccgtaatt 2820
tgcagttaat attaaactcc tgtggacgaa gctaccgctc aaacgccact tgccccgaa 2880
cctaagaaca gaatatgtat ggcactaatt accatattac ttctatcat ggtgtttgtt 2940

SUBSTITUTE SHEET (RULE 26)

WEST

17/33

tgtttgctctg tgtttaaagt taaggatggt atacccttcg tgctgctac atatatatag 3000
tgggcactat aatattacca ataaattaac gtccatatat aagaataata ataaataaat 3060
aaatatttct atacaataa aggttaagta atgttggtgt tctcgtggat ggggatctta 3120
tcttctctct cgctatcttt gtttatcgta tttcagtga agttgttcaa taaaagtcct 3180
ttgttcaaca agtgattcct tctctctctg tctttctttt cactttcgta ttttctttag 3240
gtataagggt gcaaaaatag acaggaatat cgatcttggt ataaaattaa aatcggtttg 3300
ctgatgtttt aattagttag aaaaaagaag acatatattt atcgtaattc ctgttcatga 3360
ttataaga 3368

<210> 8

<211> 7235

<212> DNA

<213> Glycine max

<400> 8

gtcgactcga tctcaaattt tatttcattt aaaataaaac ataatttaac tttcgtctct 60
cttctcttatt gtatcattat aaaagtagga aaacaaatat aaattagaac aaacataata 120
ttaattaata agaataattt gtttggtgct ttgaattttc tattctaata acattaggta 180
gtaataaaat taagttgagt ttcatttttt tgaaagaatt aacttaataa ttgtatattt 240
ttgtttaagt ataatatattt agataatgta ttatcacatt aaaaatttag agtgatagac 300

SUBSTITUTE SHEET (RULE 26)

WEST

18/33

aaattatgtt tattaatcaa tattatgttt atttaattgt ttgttttaag ttaggtttgt 360
tttcatattt tttttagtgt ttttattata atgagaaaaa aaatggagga taaaagataa 420
aaattatatt atattttact cttaaataaa acaaaatttg gagtcctaaa aattagttaa 480
atgaaaactt gttgacatag gtctaacta ttcaataatc atgttagctt atttgtgttc 540
ttggactcct tcattaacgg taatagatga atgaatttat tgcatttctt tcacttttga 600
ttactaaaat ttaaattttc atttttcgaa aatgaattcg ttactatttt atacatttaa 660
aaataaaaat gattatttca tcttataatt atcaaataat atgcctgtga tgatgatttt 720
tttcaaaaat tgaaaatgtc ttattgccta gttaggctat agaacttttt tggctcatct 780
caatcgcatg ggcacatggt taccgcgttc attgggtaat tatatttatg tatttaaatt 840
aattctatta aaaccaattc tattaaagct gaaaactaaa atgacactta atttctgtaa 900
gagtcgtgta attagcgagc cagagtaatg caaccaaaga gcttctttcc ccaccttaat 960
tcctttataa tgaatttgat tgataccata catagctaag cttttttttc ttctgattta 1020
catagccaag ctttagctgc tatatataat atgttggtta tataattttg acatgttgca 1080
ttacatgtta ctaactggtt gcaacctcca gttttggtta tcgaatatat gctggccggc 1140
accaactaca aaaaatttct gtatggctgg ctactaatac tatatcttca aatgtctcac 1200
tttacatgga ctgagtgcac aacgggcaat tgcagtgtc ataaggaata cgcatgcaat 1260
cattttatca tcgcagattg ttgcttttcc ttactgtaa aaaaaataa aaataacaat 1320

SUBSTITUTE SHEET (RULE 26)

WEST

19/33

taattgagca ccgaatgaac tagaccgatt gcatgcatta taatacatga agaaaattat 1380

ttatactaata cttttgtttc aagttttaaa gttgactttt ttgttttaca agcacaataa 1440

caatcaaata accaataatt taatcattaa aaagataaac aaactcatta gataaaataa 1500

taattatgat aaaatatgtt taagtcctct atatttaagt tgaattttgt ttttagtctt 1560

tcaaattgtg catttttagtc tctgaaattt tattttaaaa aaatagtctt tgggcttact 1620

ctccaaaatt tgcacatttt aatttttaggt tattgattaa agattaaaaa catttttttt 1680

tatcaaattt agggaccaaa ttacttgata taaatactca agaactaaaa acaaatttca 1740

ataaaaatat aagggattaa aaatatattt tattctgata actaacaatc atcaaatttt 1800

atgacaatca tatgtaacct ttataatat aattgtaaga ataattaatt ttaataaata 1860

aataaataaa tgagacttat aatagtaa atcatatgtt gcctaggtaa taacaatgat 1920

aacaagtaat agaattatca ttcttattgt atctattaat taatatatta ttagatgtat 1980

taattagtat atatattttt tactataaat agtatatatt attaattatt atacatttat 2040

taaagtatat tcattaaaga ttttgaggga ggggagctga agccactaac ccccgtaa at 2100

ccgtccttgc aaaaaacag acatgagaat ggttttgtat actccacaat ttaatatcca 2160

ataaaataat ttctttttta ttttaattag gaaactccaa gattgcttta acttatataa 2220

aatctgaata acacaaaaaa ataaaaaatc ggaattaaat gttgccgat tgttgactaa 2280

ettatacaa atcttattta aatgcttaaa atcgtgtcat aatataatga atatatttt 2340

SUBSTITUTE SHEET (RULE 26)

WEST

20/33

gcaaatatat atttattata taattgcaaa tatatatctt aattttgagt ataaataaca 2400
gcatgtgagg gtgcagcaaa acacacactg agtgcaacaa agttttaaca tgaaaggcaa 2460
taatacactt ttgttgcaat tattctacac tactctcttc ctgtttcttg tagtgtcaag 2520
ttcatcttca acagggaaatg aaagtaacga tgacactaac agtaaagaag tttatatcgt 2580
gtacatggga gctgcagatt caacaaaagc ttctcttaaa aatgagcacg ctcagattct 2640
gaattcagt ctaagaaggt acgtataatt acataatatt attattatat gggcaccaat 2700
taattaattt gatgattgat gtgtttacat attttgtgtg aatgaattga aggaatgaga 2760
atgccctagt acggaactac aagcatgggt tctctgggtt cgcagctcgt ctatcaaaag 2820
aggaggcaaa ctcaattgct cagaaacctg gtgtgggtgc tgttttcctt gacccattc 2880
tgaagctcca cactacacgt tcatgggatt tctcaaaag ccaaactcgt gtcaatateg 2940
acaccaaacc aaatacctg tccgggtctt cttttctctt atcagacgtc attcttggtg 3000
tcttagacac aggttggtcca taatcaaaaa aaaaaaaaaa acatgatata tatgtgtgtg 3060
tttcattttt taaaaatgtt aataataata tatacaaaaa tggaatattt caggcatatg 3120
gccagaggcg gcgagtttta gcgacaaggg ttccggctct gttccatccc gatggaaagg 3180
cacctgcatg acatcaaaag acttcaattc ctcttgttgt aacaggtaaa ctaaaatgtg 3240
aaaccataat aataataata ataataataa taaatatata aaggcgaacg ttattaatta 3300
ttaattatta ttagaaaaaa ggtgatttca gcttgctgtt taagaagggt tggaatgaat 3360

SUBSTITUTE SHEET (RULE 26)

WEST

21/33

cctattttaat taggtagtgg atggaataac ggtaggttt gtatttataag gaagataatt 3420

ggcgcgaggt ttaccctaa ccagaggag aaaacggcaa gggatttcaa cggacatggg 3480

actcacgttt cgtcgacggc agtgggcgtg ccggtgagtg gcgcacgtt ctatggtctg 3540

ggcgcgggga cggcaagggg tgggtccctt gagtcaaggt tggcggttta caaagtgtgt 3600

ggggcttttg ggtcatgtcc tgggtcggcc attcttgctg ggtttgacga tgccattcac 3660

gacggagtgg atatcttgtc gctgtcgttc ggtggattcg gtggaaccaa aaccgatttg 3720

accaccgacc cgattgcgat tggagcattc cactccgtcc agcgcgcat cctggtggtc 3780

tgcgccgccc ggaacgacgg agaaccattc accgttctca acgacgcacc ttggatttta 3840

accgttgacg cttccaccat cgaccgtgat cttcaatccg acgtggtctt gggtaataac 3900

caagtcgtca aggtacctac atattctact ttaaatcggg gcagtgcac taatgtcatc 3960

ttttctcatc gttgataatt attaaacttc aggaagagc cataaatttc tccccctttt 4020

taaattctcc cgattatcca atgatatatg ctgagtctgc tgccagggca aatatctcca 4080

acataactga tgcaaggtag gtactctaaa aaccatttgt cgtttcgtat tggacaaaact 4140

tcaaatcaag caatcaacta agcaataaca aacaagtgtt tcatacacia ttatatgtaa 4200

tactcatata taacctctta gcaaatgatt aaatcatttg tcacatgcag acaatgccac 4260

ccagattcat tagatccaaa aaaagtcata gggaagattg tggtttgtga tggaaaaaat 4320

gacatttatt attcaactga tgagaaaatt gtcatagtga aggcgttggg aggaataggt 4380

SUBSTITUTE SHEET (RULE 26)

WEST

22/33

ctgggttcata ttactgatca atctggatca gtagcatttt attatgtgga cttcccagta 4440

acagaggtaa aatcaaaaca tggcgacgca atcctccagt acatcaactc aactaggtaa 4500

ggatattata tagcacttga aagaagcaac attcttgatt aattttagaa ttgctttga 4560

tcacgagtta ttttctttta attcttttgat catatatgta atataaagcc atccagtggg 4620

aacaatacta gcaacagtta caattctga ttataagcct gctccccggg tgggttattt 4680

ttcatcaaga gggccttcat tgattacaag caatgttctc aaggatgat atgacgatcg 4740

atagaattat acatatcaat catcatctc aatatgctca ttgctcaaac actaaacaga 4800

acattcatc tttttttctt tctttcttct tagcctgata ttgcagcccc gggagttaac 4860

attctcgctg catgggttgg aatgacaca tcagagggtc caaaaggaag aaagccctca 4920

ctatatcgca tactctcagg aacttccatg gctactccac atgtttcagg gcttgcatgc 4980

agtggtcaaaa gaaaaaacc cacttgaggt gctccgcaa tcaaactctg catcatgact 5040

tcaggtcacc catttgataa tgtgatctaa gtaagtaatg tgatccagca aaatgtacca 5100

taccaactca tatcattcta taaattaata tgtatgcagc aattcaaaat gacaatttga 5160

aggggtcccat aacaacggat tcaggggtga tagccacacc ttatgactat ggagcagggg 5220

caattacaac atctgaacca ttgcaaccgg ggetagttaa tgaaaccaac aacgttgact 5280

acttgaacta tttgtgttac aatggactta acataaccat gataaaggct atctccggaa 5340

ctgtccccga gaatttcaat tgtcccagg attcgagctc tgatctcatc tccagcatca 5400

SUBSTITUTE SHEET (RULE 26)

WEST

23/33

actacccttc catagcagta aacttcactg gcaaagcaga cgcggtcgtg agtagaactg 5460

tcacaaacgt tgacgaagaa gatgaaacag tgtacttccc cgttgttgaa gtccttagtg 5520

aagtaattgt cacactcttt ccatataatc ttgagtttac gacaagtatt aaaaaacaaa 5580

gctacaatat tactttcaga ccgaagacct ccttgaagaa agatttgttt ggatctatca 5640

cttggagtaa cgacaaatat atgggttcgaa ttccttttgt attaactaaa tagtgaaatt 5700

aaaaagtagc gatgaataaa tgcaagctaa gttcttcgtg gtgcctacac tcgagtcctg 5760

attatttatt attcatatgc cttctgtttt aatttaattt attatacttt cagcctctct 5820

aatatgtttt tttcttttgc aaatatataa gctgacttac tatttacct caaaattagt 5880

tccaacttat tcactagccg ttgcccctca gcttaattaa aaaaaagaaa tgtgatttaa 5940

ttacattaat tatactgga tcgtagtaac ctccgatttt tacacgggtt ggtaattcaa 6000

catcaatttc atgcttcaaa tgcaaacctc tcaaaagtag ttgcagacta aaatgatgaa 6060

ttttaacaa aacttgtaaa aaggtaagg ggaactaggg aagtgagctc ataaaataag 6120

gaaactgttt cgactgagtt ttgagtaagg tgtggctgaa ttttggttg agttgtggtg 6180

agctgtagca gagtttcgac tttgtgtggt taagctatga cttagttttg acgaattgtg 6240

gtgagttgtg gtogaatgga atcttggatc tctaataccg gtgtaggaga agtacctaca 6300

aaaaggactc caacaatcaa ctcaattgga tccgagatac ttatgtatcg atgtatgaaa 6360

tgattaaaac ataccttggt gtgttttatt tatgtcaata tatacatatt cgacattaag 6420

SUBSTITUTE SHEET (RULE 26)

WEST

24/33

gaggttacac taactagata gtcctctaag cgttgctga tcgtgaaaag ttagtgatca 6480
tcactatcga gttcataggt tgtacggtac ataaatccca ctaatcgagt aactatcaag 6540
ttcatggggg ttgtacggtg cataagtctg tcagattccc atgatgggta ttgctaagtt 6600
gaataaatcg ggcataatca ttacacgagt ttaagatgat ttaattttcc ttatatatca 6660
ttattttata ctgggtgagt gttttctttg aaaaactgag gtgtggatcc aacctcttga 6720
gaagtgcctt taaaaaatg aggtttggat cctatctctt gagagggtgct tgaaaaagct 6780
tatcaaaata gttattggca ttcaatgttg ttttcaaccg agggcaatta aacacacctt 6840
tgattagtgg gcaattgaag ttgaaagaa gcttataaag gatagtatta ttatactatt 6900
acaaccaagc aaacaatggt tacttcaaag ttgatacctt attgatataa tgtattattt 6960
actatgagtt aaaatgtaat ttgaaaaaaa aaatgtaact gaaaatgttc ggatgatattt 7020
catattttgt aaaaaaaaaa aaaaaagaaa aaaaagtatt taggctaatt tttgcaaagt 7080
acattttggt atgaaaaacc aaaaacagaa gtaatgcatt ttgtgccatg gcagaatgca 7140
gagtatctac ccaggattca ttatgaacaa cttaatgcta agcatcatca gttaaccccc 7200
caaattaatc ttgaaagcta caattctatt ctaga 7235

<210> 9

<211> 8310

<212> DNA

<213> Glycine max

<400> 9

SUBSTITUTE SHEET (RULE 26)

WEST

25/33

gagctccgcg aaatttggtta tggccatact cttccttgcg agccctcttg gtctcttggt 60
caagggtctct tgcggtagtt gcattctctt cccgtaattt ggcacaactcc ttccggatgt 120
gtgtagcggc taacttgaac ttctccttgg caagtttcgc ctttcctaac tcgtttttga 180
gagcttggac ttcttcgtct tcttcgggtg cttcgaaact gtctttgctg acgactttta 240
acttggcgag ccaatctaaa cctcgtattt gaactttcag ccattcatga taaccaccaa 300
tgatgccatt acgaatgcc ctaagttctt gatctttcct taacgggggt tcccatgcct 360
tatggattct ttgtatagcc ttgaaatttt gcctgcgaa atctctcaca aggaaaggag 420
aaatcctttc ttccatcggt gtccctca tggggtaacc tagttgtctt atagcgagcg 480
cggtattgta gttgatacaa cccctcgttc ttatcagtgg aatgtttggg taccctccac 540
atgagaaaag gactccctcc ttctcttctt tccatcgggg gaaccaacta attgttctac 600
ctctatctc ggccaagagt tggcccatt ctattctct cttttcagca cagagtgat 660
ggctttggag cggacatgga tgcctcgtgt ttgctggaa cagggtgaa accaaccaaa 720
cacagagagc gggcaagcaa cagatgatec atgcgtact cttctcgac cttcgggtcaa 780
atgtgtcaaa taaatctgcc aagacagcta ccaccgggt ttctgtttg gaaagaggac 840
gaccccaaag attagcaatg ctaacacatc cataaacggg acccaatctt cttgatttgc 900
catatccctc gccttgctt ctaggtactt ccgtggtagg cccgctatgc cgtttcgagt 960
ttgttttacg cggtcctaac ctcttgctga atctttgacc acggtcgaa ttctgctcaa 1020

SUBSTITUTE SHEET (RULE 26)

WEST

26/33

agaggggaga caccgggaga aaagatatga ttttcttccc ccgagaggac atcctagaat 1080
ctcctcaa at tcttcaatgg tcggtaccaa ttggaagtct ccgaacgtga agcatctcaa 1140
aggctgggtcg tagtattggg tgagtgaagc aatggcctct atggatacct ctggtatggg 1200
caaatactaa atcgatcttg gattgttggt gggtttgttg aagtattttc taaccttttg 1260
aatatgttac taatgtcaaa tttattatgt gttataaaaat ctttttgtct gggtttattt 1320
tctaggggtt tctttaattt ttccagaaaa actttctttt cctggggatc aggtatagaa 1380
ttgattgcct caaacaagag atcgatctct ccgaaattgc acacagtggg caatgtctca 1440
gcatactctg tttatagttg tggattctat aagtcaattt agattattgt taattaccta 1500
agttattgtt ttaaattcat acataattaa ctttgtctta acaacaacag acatctgact 1560
gagacaagta tgtgagcggg gaattctgat gtgtatggat tcctcgagcc acagtctatt 1620
cagagatctg ggcaatcgca gtttgaatca gaaagttaca tcaagagttg gatgcagagt 1680
tcaaaacgag atgtctacct tggagcctac ctgaatgggt aagtcaaata aaacaactaa 1740
atttaataa tatataatac tacaataacc catattcatc tccactgcag cggacacagg 1800
aaaatggtct ttattctgcc taaggaaaac cttgttgtct gggtttgttc cttgcataac 1860
aggtcagaca actaccttaa gggaataatt aacagggtcag tgttggtttt aatacatgtg 1920
cattagcata actcaacaac atcaatattt taatgttctc cgtattcaac agtgctttga 1980
aaggacttga tgatactcca caatctaaat ccaaggctgg tgctaggtgg atcgttggtta 2040

SUBSTITUTE SHEET (RULE 26)

WEST

27/33

aggtacgtga tttaaataaa acttctactt atatatactg cttatgtgtg tgtacactaa 2100
ttgtagttaa cgtttaatta atatccaaat ttcattatgt atttagtgta atagacaaaa 2160
aggaagcact gagtgcggct attacatcat acactagatg tccatgataa tcttaggaac 2220
ttttaggaat aattgggaaa cegtaattgt ttatttcaaa caaagtttat tttcttgtca 2280
ttggtattac attattaact tatgttttat ttgatcatgc agtattttaa cgatgttaaa 2340
ccattggaag caaagagatt gaagggtgctt cgcacccagt gggcacaatt ttatctcaaa 2400
gttacaatac aaagtttaga tgttagggaa ctatttcatt ttaggttact taattagttt 2460
actaccttgt ttacattttt tgccaattgc tatgccattt gaacattgaa tattcttaat 2520
tgatagttat taataaatca ttattcata gttatcaatt gatagtttac tatagcttta 2580
tactaaaaac cgtttaaag cagaatgcaa atgatatatg ctgtgatcta tggctcgatt 2640
tcaattttac aggttcaatt ttgagttttt ttgtaaaaac aaaaagtgtt ttaaaaaaaa 2700
ctaaaaacaa aatcggtttt ttacaaaaat cgatgttaac atacaactta acatcggttt 2760
ctcaaaaaac cgatgtttgt gtatattaac attgattttt gtaaaaaacc aatattaacc 2820
tatgaaaatt taacattggt tttatacaa aaccgatgtt aacttataga ttaacatcag 2880
ttttgtataa aaacttatgt taacgtttct aacttaacat caatttttat agaaaaccga 2940
tgtaatatata tgagttaaca tcgattttga tacaaaactg atgttaatct ataagctaac 3000
atcggttttg tagataaccg atattttaaa tttacgtta aactgatgtt aacgataata 3060

SUBSTITUTE SHEET (RULE 26)

WEST

28/33

ctttcaacat cgattaaaaa tcgatgtaga aagtcgtaaa taaccgatgt agaaaatcta 3120
ttttctaata gtgtccatct caatatatgt ttggatagtg tcggcattac cctggtttga 3180
ctaagtcttt tgcactcaaa ttgtcacacc ataaaatcgg tgttctgaga atgggcaagc 3240
taatcttctt gatccattga gaacctaat ttcattatct tttaccta acataaaaaac 3300
acaaaaatat aagaaaataa aataaaatcc ttatctaat tcagaacata aaataaataa 3360
catctttcat gttaatattt taaaattcat tattttgaag ccaaattggg ttaaaattta 3420
taacaagtga cttaaaatta aaatttccaa aaataagtag aaaattagat aatttaaaaa 3480
ttaccaaact cattttatcc ttcagtactg ttatgtactg atgcagtgt attaccaca 3540
ttataaatgg taattacaaa ctagtactac cttataaatt aattacccc attctctat 3600
cttacaata aaattaaaaa tacggacaaa catggcagtg gtgtggtttc gcttaaaaa 3660
taaaaatata caaacatgaa agtgatgtgt tttcgttgtt atatataa atgctcacga 3720
tcgagactca actaaatgat caccgaaatg ttttcttag gatgaagcct gttgctaatt 3780
tcttcattct aaaaatcggc cattacgcaa tttatagatt tgcgagatt ttggttctt 3840
ctttaaattt ggcataaaa acttatcaat gtatacctta tggctgcatg aaattaccta 3900
ttcttatggt tgcataata atgttcatac tttttatatt atttattta gattaaattg 3960
taattttgat cctattttt aaattaaggc attccattaa tggaccattt atgtgacaaa 4020
agattaaata acttaaaaat acaaaaaggc cttagtataa attaaattta aaatatttta 4080

SUBSTITUTE SHEET (RULE 26)

WEST

29/33

tataaagata aaataataaa accataaaat acttacttta tttagttaat tttataaata 4140
ctattttaca tgtatcatta attagtccag aattattaat ttttttaatt ataaaaattt 4200
ataaattgaa ataaaaatatt taatatttaa atatatttaa catttaattt tacatgtact 4260
tatttttctt ataaaaattt attttaaaat aaatcaacta aaatttggtg ttattttagt 4320
ataattttaa aaaaatacat aaactaatat atagattatt ttaaaattat tttatctaata 4380
ttatgaaaat taagtagatt taaaataaca tttaactaat ctaaaattta aaatttatta 4440
cttgttataa ttaaaataaa ctttttcatt atttatatat aaaaataaat gtagataatt 4500
ttatcaatta cataaaataa tttatatata actaatttac atattaattt atttaatttt 4560
cctcatttat ataattgata ttgaaaaact tatttactaa atatttttta tagttaaaat 4620
aaaaatatta acatgttttt ttcttaaaaa cattaaatat tcaaatttta ttcatttatt 4680
taattaaaat aaaggttaatt tacatattaa aataaattac ataatttgta taatttatat 4740
attaataaaa aaatatataa attaatttta attagtcaaa taaaaataaa aatatataaa 4800
ctattcaata tatatatata tatatatata tatatatata tatatatata taatataatg 4860
acatacaaga aaattaaaag gatttataaa aaggatttat atattaaata ttttttaatt 4920
cataaatttt tataatttaa aaaattaact catacataaa taatatttaa aaaattagtt 4980
aaaaaataa atatcttaat agtttggtat tttaatttta aataagagat tataagttta 5040
attcttatta ttttttttca taattttaac tcttaccaag tatttttttc actttgttta 5100

SUBSTITUTE SHEET (RULE 26)

WEST

30/33

atctctattc tctctctctt tatatatata tatatatata tatataataa atttctaate 5160
aactaaatag tgtatatattt acatgcagag tacataaaaa ttaaaattgg gtataaatagt 5220
tattgtaagt catctcactc actcttcttc ttagagtcta tatatatgca atatctctgg 5280
tctccctatc accttctctt ctctaagcac ttacttttt ttttcagcca tggagtttca 5340
ttgccttcca atatttcttt atctcaatgt gagcataaac ctctcttttt catctcgttt 5400
tttaacctgt attgcgtgaa gttggatttc ttcattaatt atcattttta ttatagcaat 5460
aagtatcaag tgatgggttt ttgcatgaat tatgtagttg atgttgatga cagccaatgc 5520
tgcgtaact cctagacatt actgggaaac gatgcttcca agaactccct tgccgaaagc 5580
aatcacagag ctactaagcc ttggtgagta agaaatcaaa ttgaaataaa ataagcaaca 5640
cttttgtaat taaatctgaa acgataaatg tacgtaaaca aaaaatagaa ttagaaacca 5700
atgcaaacag ctataagcct ttttttgtaa tagctaattt gccttaatta aaccattgaa 5760
tgactcaata tgttcattct ttttccctt tagtagaata ctgcccttct tcatattoga 5820
tatactattg gcttccctag tgaattctta ttctaattgg tgcataagctg tgttctcact 5880
aagcctcctt tcccatgtct ttgattctat gaaacgaaac tttcaaaact tggaatcacc 5940
tattaagtac aataaagaaa taattaattt ttttttata cttaattat tttatgaatt 6000
tggttccctt attttatttg atatattttt tacaaatttg gtttctctat tttttaatt 6060
gtttattttt tgttcacttc ttaacttaat atccaacatt tcataaatat gctgacataa 6120

SUBSTITUTE SHEET (RULE 26)

WEST

31/33

tactagatta ttatatcacc tgtggggata ttcatgttgc acaattagtg gaggctcaaa 6180
ttcgtgaaaa aataaataaa atttggagag accatattca ccaaatacaga atagtatata 6240
gagacaaaaa aaaaaaaaaa aaacaaatta tgcaacgaat gaaaagttaa ccattatatt 6300
tagtattgag taatttgaag tcagtgtgag acatggcaac gtagttaact tactttaatt 6360
aaaaatatga ataattaaaa taggaatata taacggattc ggattctctc cagcaatgat 6420
gaggacgtag gattaatttt atggctgaga gtaaatccaa aaagatacgc aaaccacgta 6480
tcaccgctac tgcttgacca aatggaagcg acttcaaacc aaaattagga caaagaaaat 6540
aaaaatacaa atgtcaataa actcaatata caaagacaga gtgtaagttt tactaaatgg 6600
aaagctggaa ttgaagtaaa agctttacat atatattgct tgctaaacaa ttttgaattt 6660
tttttatctt ggattcttgc taaacaatta attatatcta ttaatatctc totaataata 6720
tgggatacac ttattattat tattattatt attattatta ttattattat tatttgattt 6780
aatataatt ttgatatttg actaatatta aatttatttt aaacaatcac aaatttcgaa 6840
taattaatga taatattaag ttattttcat atttgactag tatcaaaaag ttatttacac 6900
atttgactta ctgttatttt attttagttc tctagttttt aaaagtataa taagttagtt 6960
ctttcatttc ttttctattt tccattttta ttaaaaaaaaa ttcaattttt attttatttt 7020
gtaacaatgg gctgaaaata gctactaaaa ttttggctgt gtttttcttt accaatcaga 7080
aagtaggtcc atatttgaat atgcggggaa tgatgaccag tcagaaagta ggtecatatt 7140

SUBSTITUTE SHEET (RULE 26)

WEST

32/33

aggatacgtt ggctataatc aagacgagga tgatgtgagc aaacacaata tacaatctt 7200
caacaggttg tttttcttgg aagaggacct gcgtgctggc aaaatattca acatgaagtt 7260
cgtaacaac aaaaaagcca cagtcccggt gctaccgggc caaatcttga acaaatacc 7320
gttctcagaa gataaaaaga agcaagtgtt ggcgatgctt ggcgtggaag cgaactcaag 7380
caacgccaag atcatagcgg agaccattgg tctttgcca gagcctgcaa cggagggaga 7440
aaggaaacac tgcgcgactt cgttgaggct catggttgat ttctgctttt ccgcgctcgg 7500
gaagaacgtt ggtgctttct caacagagaa agaaaggga actgagtctg gaaagtttgt 7560
agtggtagaa aatgggggtga ggaagttggg agatgataag gttattgcct gtcaccaat 7620
gagttaccct tatgttgtgt ttgggtgtca tctagtcca aggagtagcg ggtatttggt 7680
gcgcttgaag ggagaagatg gggttcgagt gaaagcagtt gttgcgtgcc acagagacac 7740
gtcaaagtgg gaccataatc atggggcatt caaagtgtc aatcttaagc ctgggaatgg 7800
tacagtatgc catgtcttca ctgaggggaa tcttcttgg ctccaaatt agattaatta 7860
ccatatacat atttgtcctt gttctatcct taaataagt gaatcacctg agaattgtg 7920
cgtaatgagt tgtttgtctt tgtggaaatt gttatctgtc ttgcatcacc aaataggtat 7980
atataaaata acaggagcgt ggtatttggt gcacaaaaat ggatttcaac cgatcaaaaa 8040
aatatagcct ttaccaatta gaagggttg gctttgttag caaataataa aaataaaata 8100
tcttgatgga taaatggttg ctaagttgat taagattgtg gcagaatacc aagtcaatga 8160

SUBSTITUTE SHEET (RULE 26)

WEST

33/33

atagtcctc acaagcatca aaagaaaaga taatgattcc ttgaaaatag aaagcacttt 8220

gtgttttgaa ttcaaaatgc aactggaga gttgttgag gttacaagcc agatcgagtc 8280

gactcccttt agtgagggtt aattgagctc 8310

SUBSTITUTE SHEET (RULE 26)

WEST